

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-----------------------------|--------------------|----------------|-----------------|---|--|--|
| Acid Fast Stain (for bacteria) | 8.50 /sample | Bacteriology | M-F | 1-2 days | 1) Affected tissue, lesion, or fluid; 2) Impression smear | 1) Amies transport media (swab) or sealed sterile container; 2) slide mailer | 1) Refrigerate. Ship in insulated container with ice pack. 2) Not required |
| For the direct microscopic detection of Mycobacterium species. *May qualify for New York State contract pricing | | | | | | | |
| Acidic-Extracted Drugs | 70.00 STAT+25 | Toxicology | M-F | 2 business days | 1) 4 ml heparinized (preferred) or EDTA plasma or serum; OR 2) 10 ml heparinized (preferred) or EDTA whole blood; OR 3) 50 ml urine | Heparinized tube(s), no gel clot; or red top tube with transferred heparinized plasma, labeled accordingly. Less ideal: red top tube with transferred serum or EDTA plasma OR red top/lavender top tube(s) with whole blood OR urine in leak-proof container | 1-3) Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Qualitative drug screening and confirmation. Analytes include, but are not limited to: Acetaminophen, Barbiturates, Bumetanide, Chlorothiazide, Diclofenac, Diflunisal, Firocoxib, Flufenamic Acid, Flunixin, Flurbiprofen, Furosemide, Griseofulvin, Ibuprofen, Indomethacin, Ketoprofen, Meclofenamic Acid, Meprobamate, Methocarbamol, Naproxen, Oxyphenbutazone, Phenylbutazone, Nabumetone, Salicylic Acid, Sulfa Drugs (Sul diazine, Sulfadimethoxine, Sulfadimidine, Sulfafurazole, Sulfamerazine, Sulfamethoxazole, Sulfanilamide, Sulfapyridine, Sulfathiazole), Sulindac, Thiosalicylic Acid, Tolmetin, Zomepirac. | | | | | | | |
| Aerobic Bacterial Culture | 40.00 /sample | Bacteriology | M-F | 2-7 days | Affected tissue, lesion, fluid aspirate or swab in bacterial transport media | Amies transport media (swab) or sealed sterile container | Refrigerate. Ship in insulated container with ice pack. |
| For the isolation of aerobic bacteria. Tissues are not routinely pooled for aerobic cultures. When multiple tissues are submitted and aerobic culture is requested, we will culture each separately, and a fee for each tissue will accrue. To detect Salmonella from GI samples, a Salmonella culture should be requested. Salmonella is readily identified in aerobic cultures from non-GI sources. To detect Listeria in brain or CSF specimens, a Listeria culture should be requested. Listeria is readily identified in aerobic cultures from systemic samples of septicemic animals or fetal tissues. Streptococcus equi is identified in routine aerobic cultures. If no other aerobic bacteria are of diagnostic concern, a Strep. equi culture can be used to rule in/out Strep. equi only. For Campylobacter spp., Mycoplasma spp., Ureaplasma spp., Yersinia spp. and Brucella spp., individual cultures specific to the bacteria should be requested, as they are not detected reliably in routine aerobic cultures. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not included in the testing fee. Susceptibilities will be performed only if specifically requested. See Antimicrobial Susceptibility (MIC) and Antimicrobial Susceptibility, Ophthalmology. Preliminary results will be provided one day after sample processing begins. | | | | | | | |
| Alanine Aminotransferase (ALT) | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| See Eclinpath for more information | | | | | | | |
| Albumin | 8.50 | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship in insulated container with ice pack. |
| See Eclinpath for more information | | | | | | | |
| Alkaline Phosphatase (ALP) | 8.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| See Eclinpath for more information | | | | | | | |
| Alkaloid Screen | 331.00 /sample | Toxicology | M-F, as needed | 5-7 days | 1) liver; 2) heparin whole blood; 3) urine; 4) other materials | 1-3) unbreakable sterile container; 4) Heparinized tube, no gel clot | 1-4) Refrigerate. Ship in insulated container with ice pack. |
| Qualitative screening for alkaloids such as strychnine, brucine, caffeine, theobromine, theophylline, nicotine, atropine, cocaine, morphine, codeine, ephedrine, reserpine, quinine, and solanidine, on request. Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | | | | | | | |
| Ammonia | 54.00 | Clinical Pathology | M-F | 1 day | Separated plasma from a purple top (EDTA) or green top (heparin) tube | Red top (with separated plasma) or purple top (whole blood) | Refrigerate: 4 C or frozen (on dry ice) – see Appendix A for special handling instructions |
| Used to identify hyperammonemia in cases with neurologic symptoms due to hepatic failure (hepatic encephalopathy), enteric disease (colonic disorders in horses) and inherited disorders of ammonia metabolism. Call the lab at 607-253-3255 prior to drawing sample. See Eclinpath for more information | | | | | | | |
| Amylase | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship in insulated container with ice pack. |
| See Eclinpath for more information | | | | | | | |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 1 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|--------------------|-----------------------|---------------|-----------------|---|--|--|
| Amylase, urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer pack. |
| Anabolic Steroids in Urine | 441.00 /sample | Toxicology | M-F as needed | 7 business days | urine or plasma | sterile container | Refrigerate. Ship in insulated container with ice pack. |
| Qualitative screening for boldenone, dianabol, and stanozolol. Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | | | | | | | |
| Anaerobic Bacterial Culture | 45.00 /sample | Bacteriology | M-F | 2-7 days | 1) Fluid aspirate inoculated into anaerobic transport media vials for liquid specimens(Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM); 2) Fresh tissue (>2-3 cm in diameter, minimum size); 3) Ligated loop of intestine; 4) Swab or small tissue collected and immediately placed in anaerobic transport media tube for swabs(Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM) | 1,4) Anaerobic transport media (Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM): tube or vial form - see sample info above; 2-3) tissues (>2-3 cm in diameter, minimum size) in sealed sterile container | 1) DO NOT Refrigerate anaerobic transport media tubes or vials; 2-3) Tissues (>2-3 cm in diameter, minimum size) are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection. |
| For the isolation of anaerobic bacteria. Ship overnight. Antimicrobial susceptibilities are not applicable. Please consider using the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine diarrhea submissions. Tissues (>2-3 cm in diameter, minimum size or ligated loops of bowel) are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection. DO NOT Refrigerate anaerobic transport media tubes or vials Despite labeling to the contrary, culture systems containing Amies transport media or Stuart transport media are inappropriate as transport systems for anaerobic culturing. Swabs transported without any transport media are unsuitable for both aerobic and anaerobic culture. | | | | | | | |
| Anaplasma phagocytophilum (Ehrlichia equi) PCR | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) EDTA blood, buffy coat (min. 2 mL); 2) tick: see comment; 3) tissue (min. 5 g) | unbreakable, leak-proof container | Refrigerate |
| PCR to detect Anaplasma phagocytophilum EDTA blood is the preferred sample from a sick animal. Ticks must first be submitted for parasite ID to determine if they are a vector species; see Parasite Identification test. | | | | | | | |
| Anesthetic Screen | 331.00 per /sample | Toxicology | M-F as needed | 3 business days | 1) urine; 2) heparin whole blood; 3) liver | 1) sterile, leak-proof container; 2) Heparinized tube; 3) sterile, leak-proof container | 1-3) Refrigerate. Ship in insulated container with ice pack. |
| Qualitative screening for acepromazine, barbiturates, ketamine, morphine, xylazine, tiletamine/zolazepam. Special request only; contact Lab. | | | | | | | |
| Anticoag/Rodenticide Screen, liquid | 165.50 /sample | Toxicology | M-F | 3 business days | 1) heparin whole blood; 2) urine | 1) Heparinized tube; 2) sterile container | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| Screening for brodifacoum, bromadiolone, chlorophacinone, dicoumarol, difenacoum, diphacinone, flocoumafen, warfarin For more information, see Strychnine, liquids/fluids. | | | | | | | |
| Anticoagulant/Rodenticide Screen, solids | 165.50 /sample | Toxicology | M-F | 5 business days | 1) liver; 2) stomach contents; 3) feed | 1-3) sterile container | 1-2) Freeze. Ship in insulated containers with ice packs. |
| Qualitative testing for Brodifacoum, Bromadiolone, Chlorophacinone, Dicoumarol, Difenacoum, Diphacinone, Flocoumafen, Warfarin For more information, see Strychnine, solids. | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|-------------------|-------------------------|-----------|-----------------|---|---|---|
| Antimicrobial Susceptibility (MIC) | 28.00 /isolate | Bacteriology | M-F | 2-5 days | 1) Affected tissue, lesion or fluid; 2) Feces or intestinal content; 3) Bacterial isolate | 1) Amies transport media (swab) or other appropriate bacterial transport media OR 2) Exudate or aspirate fluid in sealed sterile container OR 3) Bacterial isolate on Agar slant or plate (sealed) or swab in appropriate bacterial transport media | 1-3) Refrigerate. Ship in insulated container with ice pack. |
| Includes systemic antimicrobials. Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. *May qualify for New York State contract pricing . | | | | | | | |
| Antimicrobial Susceptibility, Ophthalmology | 30.50 per isolate | Bacteriology | M-F | 2-5 days | 1) Bacterial isolate | 1) In house culture isolate 2) Agar slant or plate (sealed) | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| Includes topical ophthalmic antimicrobials. *May qualify for New York State contract pricing . Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. Special certified training is required for the shipment of some pure cultures. Contact AHDC laboratory with questions | | | | | | | |
| Antimicrobial Susceptibility, Urine | 24.50 /isolate | Bacteriology | M-F | 2-5 days | 1) Urine; 2) Bladder tissue/stones; 3) Bacterial isolate** (see Comment) | 1-2) Sterile container or Amies transport media (swab); 3) Agar slant or plate (sealed) | 1-3) Refrigerate. Ship in insulated container with ice packs. |
| Includes antimicrobials that concentrate in the bladder. *May qualify for New York State contract pricing . Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. **Shippers of bacterial isolates must be fully trained in Hazardous Materials Shipping and adhere to all applicable regulations. | | | | | | | |
| Anti-Mullerian hormone (AMH) ELISA | 65.00 | Endocrinology | Wednesday | 7 days | 0.5 mL serum | Plain red-top tube with NO additives | Freeze. Ship in insulated container with ice packs. |
| For canine and feline serum samples only. Ship on frozen cold packs using an overnight courier service. To assure reliable results serum sample should arrive either frozen or adequately chilled (i.e. partially thawed but kept near 4 C at all times). For more information, please see: Canine and Feline Reproductive Function Tests. | | | | | | | |
| Antinuclear Antibody (ANA) | 41.00 | Clinical Pathology | 1x/wk | 7 days | 1 mL separated serum | plain glass or plastic tube | Refrigerate. Ship in insulated container with ice pack. |
| ANA, dogs and cats only Ship Overnight | | | | | | | |
| Antiplasmin | 38.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate. Ship in insulated container with ice pack. |
| Measurement of antiplasmin levels in plasma Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page | | | | | | | |
| Antithrombin (AT3) | 35.75 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate. Ship in insulated container with ice pack. |
| Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page | | | | | | | |
| aPTT (Activated Partial Thromboplastin Time) | 14.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate. Ship in insulated container with ice pack. |
| Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page | | | | | | | |
| aPTT Inhibitor Screen | 14.50 | Comparative Coagulation | M-F | same day | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate. Ship in insulated container with ice pack. |
| Ship Overnight; requires aPTT on same sample; for additional info see Submission Guidelines, or Comp Coag web page | | | | | | | |
| Arsenic, blood/water/urine | 40.00 /sample | Toxicology | M-F | 3 business days | 1) whole blood; 2) urine; 3) water | 1) Heparinized or EDTA tube; 2) urine in sterile container | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| Quantitative arsenic concentration in fluid samples. Cannot analyze in plasma or serum - Need whole blood. Do not use heparin tubes (gel tubes). Whole blood should not be clotted. Contact Lab for specifics for avian and very small animals | | | | | | | |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 3 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|--------------------------------------|-------------------------|-----------|-----------------|---|--|--|
| Arsenic, tissue/feed Quantitative arsenic concentration in tissue or feed. | 35.00 /sample | Toxicology | M-F | 5 business days | 1) tissue; 2) feed | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |
| Aspartate Aminotransferase (AST) | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Aspartate Aminotransferase (AST), urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate. Ship in insulated container with ice pack. |
| Bacterial ID For the identification of pure bacterial isolates. Charges are assessed PER ISOLATE submitted. Please ensure purity before shipment. Submissions accepted from Laboratories only. Follow all biohazard shipping regulations. Please provide animal history and suspected identification of isolate if possible. | 40.00 /isolate | Bacteriology | M-F | 2-7 days | Bacterial isolate (pure culture) | Agar slant or plate (sealed) or Amies transport media (swab) | Refrigerate. Ship in insulated container with ice pack. |
| Baermann Fecal Technique A modification of the Berlese Apparatus used to retrieve nematode such as lungworm, larvae from feces. The Baermann Technique is not recommended as a primary diagnostic technique for evaluation of parasites in feces. It is not useful for those nematode larvae that do not leave the feces or other biological material, or for detection of parasite eggs or cysts that may be in the fecal sample. Some lungworm larvae are retrieved using this technique, but some larvae such as those of <i>Filaroides hirhi</i> , <i>Filaroides osleri</i> , <i>Strongyloides</i> sp. and <i>Dictyocaulus</i> sp. are better recovered using flotation techniques. Lungworms, such as <i>Eucoleus</i> (<i>Capillaria</i>) <i>aerophilus</i> , that do not produce larvae are detected only by using flotation methods as well. Any fecal sample submitted for this procedure must be freshly voided so that the sample is not contaminated by free-living nematodes. It is strongly recommended to use a fecal flotation technique when lungworms are suspected. | 25.00 | Parasitology | M-F | 2-3 days | 15g fresh fecal sample (~1/2 cup) | Plastic leakproof container | Refrigerate. Ship in insulated container with ice pack. |
| Basic-Extracted Drugs (including but not limited to): Acepromazine, Amphetamine, Apomorphine, Atropine, Benzocaine, Brompheniramine, Brucine, Bupivacaine, Caffeine, Cimetidine (Tagamet), Chlorpromazine, Cocaine, Codeine, Cyproheptadine, Dextromethorphan, Diazepam, Dihydrocodeine, Dipyrone, Doxapram, Embutramide, Ephedrine, Griseofulvin, Hordenine, Hydrocodone, Ketamine, Lidocaine, Mepivacaine, Meprobamate, Methocarbamol, Methylphenidate, Methamphetamine, Metoclopramide, Pentazocine, Phenacetin, Phenothiazine, Procaine, Promazine, Propoxyphene, Pyrilamine, Rantlidine (Zantac), Strychnine, Tetracaine, Theobromine, Theophylline, Trimethoprim, Tropicamide, Xylazine For more information, see Appendix B. \$25.00 will be added for STAT processing. | 70.00 /sample; plus \$25 for STAT | Toxicology | M-F | 2 days | 1) 4 ml heparinized (preferred) or EDTA plasma or serum; OR 2) 10 ml heparinized (preferred) or EDTA whole blood; OR 3) 50 ml urine | Heparinized tube(s), no gel clot; or red top tube with transferred heparinized plasma, labeled accordingly. Less ideal: red top tube with transferred serum or EDTA plasma OR red top/lavender top tube(s) with whole blood OR urine in leak-proof container | 1-3) Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Beta-Hydroxybutyrate (BHBA) See Eclinpath for more information | 13.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum, EDTA plasma, or heparinized plasma | Red top (non-anticoagulant), green top (heparin) tube, or purple top (EDTA tube) | Refrigerate. Ship overnight in insulated container with freezer pack. |
| Bethesda Unit Assay Bethesda units/mL: measure of factor inhibitory activity Ship Overnight. For additional info see Submission Guidelines or Comp Coag web page | 52.00 | Comparative Coagulation | M | 1 wk | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate. Ship in insulated container with ice pack. |
| Bicarbonate See Eclinpath for more information | 9.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Bile Acids, Fasting Panel Fasting Panel Clearly mark "Fasting" on sample. Lipemia (falsely increases) and hemolysis (falsely decreases) interfere with bile acid measurement through optical interference. Efforts should be made to minimize these (by not feeding too large a meal and by separating serum from cells as soon as possible). See Eclinpath for more information | 24.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1 - 2 mL separated serum, each sample | Red top (non-anticoagulant) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---------------|--------------------|-----------|-----------|---|---|--|
| Bile Acids, Post-Prandial Panel Post-Prandial Panel | 24.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1 - 2 mL separated serum, each sample | plain glass or plastic tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Bile Acids, Random Panel Lipemia (falsely increases) and hemolysis (falsely decreases) interfere with bile acid measurement through optical interference. Efforts should be made to minimize these (by not feeding too large a meal and by separating serum from cells as soon as possible). | 24.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1 - 2 mL separated serum, each sample | Red top (non-anticoagulant) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Bilirubin Panel (total, direct, indirect) Includes Total, direct, and indirect bilirubin. See Eclinpath for more information | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship in insulated container with ice pack. |
| Bilirubin, Total See Eclinpath for more information | 8.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Blood Culture, Aerobic For the isolation of aerobic bacteria in blood. Blood culture medium should be inoculated immediately when blood is drawn. See https://ahdc.vet.cornell.edu/docs/Blood_Culture_Technique.pdf for instructions. If the blood culture media used is not the VersaTrek EZ® Draw® REDOX 1® product, turn around time may be delayed due to inability to take full advantage of enhanced incubation characteristics of the VersaTrek® incubator system. See TREK Diagnostic Systems. The media is available from our diagnostic supply center. Submissions of 3 consecutive draws are recommended. Antimicrobial Susceptibility (MIC) are not included in the fee. See also Foal Diarrhea/Septicemia Kit. *May qualify for New York State contract pricing. | 30.00 /sample | Bacteriology | M-F | 5-10 days | Whole blood in blood culture medium | 40 ml VersaTrek EZ® Draw® REDOX 1® aerobic blood culture bottle | Do NOT Refrigerate |
| Blood Culture, Anaerobic For the isolation of anaerobic bacteria in blood. Blood culture medium should be inoculated immediately when blood is drawn. See https://ahdc.vet.cornell.edu/docs/Blood_Culture_Technique.pdf for instructions. If the blood culture media used is not the VersaTrek EZ® Draw® REDOX 2® product, turn around time may be delayed due to inability to take full advantage of enhanced incubation characteristics of the VersaTrek® incubator system. See TREK Diagnostic Systems. The media is available from our diagnostic supply center. Submissions of 3 consecutive draws are recommended. Antimicrobial Susceptibility (MIC) are not included in the fee. See also Foal Diarrhea/Septicemia Kit. *May qualify for New York State contract pricing. | 30.00 /sample | Bacteriology | M-F | 5-10 days | Whole blood | 40 ml VersaTrek EZ® Draw® REDOX 2® anaerobic blood culture bottle | Do NOT Refrigerate |
| Blood Smear Evaluation RBC (morphology, parasites), manual differential cell count, WBC, Plate smear est. Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panels For instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1) EDTA blood (> 1/2 full) AND 2) 2 dry (unfixed, unstained) blood smears | 1) lavender top tube; 2) frosted-end slides | Refrigerate blood; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment. |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|-----------------|-----------------------|---------------|----------------|---|--|--|
| Bone Marrow Cytology | 60.00 | Clinical Pathology | M - F | 1 business day | 1) At least 4-10 dry (unfixed, unstained) bone marrow smears | 1) Frosted-end slides | Hold at room temperature; DO NOT refrigerate smears; protect from chilling during shipment with insulation. |
| <p>Report includes – Specimen type and site (provided by submitter), Smear description (overall cellularity, assessment of all hematopoietic lineages, other cellular components and marrow iron by Prussian blue staining), Smear interpretation. Interpretation of bone marrow smears requires a recent CBC (within 24-48 hours). All bone marrow samples without a recent CBC should be accompanied by an EDTA blood sample with 2 blood smears and a CBC should be requested (use 'Other diagnostic tests requested' field to the right of the Clinical Summary field to request a CBC). Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing For instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear</p> | | | | | | | |
| Bordetella PCR | 36.75 | Molecular Diagnostics | M-F as needed | 3 - 5 days | pharyngeal swab; Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. | Clean screw-topped container | Refrigerate. Ship in insulated container with ice pack. |
| <p>This assay detects Bordetella DNA through DNA amplification. Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. This service is performed pursuant to an agreement with Roche Molecular Systems.</p> | | | | | | | |
| Bovine Metabolic Panel | 30.50 | Clinical Pathology | M-Sa | 1-2 days | 2 mL separated serum | plain glass or plastic tube | Refrigerate |
| <p>BHBA, NEFA, UN, Albumin, AST, Cholesterol, Triglycerides Includes NEFA, BHBA, AST, ALB, and BUN; for additional info see Clin Path panels</p> | | | | | | | |
| Bronchoalveolar Lavage Count Cytology (smear exam and count: 56.00) | | Clinical Pathology | M - F | 1 business day | 1) BAL fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate. Ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect slides from chilling/coolant by insulating separately during shipment. |
| <p>Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation, respiratory. Please also submit 2-3 freshly prepared smears of the BAL fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p> | | | | | | | |
| Bronchoalveolar Lavage Cytology (smear exam only) | 48.00 | Clinical Pathology | M - F | 1 business day | 1) Smears (at least 2-3 unstained smears preferred); and/or 2) BAL fluid in EDTA (no cell counts provided); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only. Ship in insulated container with freezer pack for next day delivery. Hold slides/smears at room temperature; protect from chilling/coolant by insulating separately during shipment |
| <p>Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation. With fluid samples of BAL, please also submit 2-3 freshly prepared smears and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p> | | | | | | | |
| Bronchodilators | 165.50 per hour | Toxicology | M-F | 1-3 days | 1) 10 mL heparinized whole blood; 2) 25 - 50 mL urine | 1) Heparinized tube, no gel clot; 2) sterile, leak-proof container | Refrigerate. Ship in insulated container with ice pack. |
| <p>(Clenbuterol, Albuterol, Turbutaline) Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis.</p> | | | | | | | |
| Brucella Culture, non-canine | 25.00 /sample | Bacteriology | M-F | 5-14 days | Affected tissue, lesion or fluid aspirate | Amies transport media (swab) or sealed sterile containers | Refrigerate. Ship in insulated container with ice pack. |
| <p>For the isolation of Brucella species only. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Refer to Brucella canis Culture for canine submissions.</p> | | | | | | | |
| Cadmium (Atomic Absorption Spectroscopy - Graphite Furnace) | 35.00 | Toxicology | M-F | 3 days | 1) 2g tissue; 2) 100g feed | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 6 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---------------------------------|--------------------------|------------------|----------------|--|---|--|
| Calcium Measures total calcium in serum, plasma and fluids. See Eclinpath for more information | 11.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Calcium, urine Measures total calcium levels in urine. | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Campylobacter jejuni Culture For the isolation of Campylobacter jejuni. Test is also part of the Enteric Panel 2 and Enteric Panel 3. Antimicrobial susceptibilities are not applicable. Recent data has shown a dramatic shift among the species of Campylobacter involved in small ruminant and camelid abortion cases. Increases have been documented in the incidence of Campylobacter jejuni abortions with research showing that it has now replaced Campylobacter fetus subsp. fetus as the predominant cause of sheep abortions in the United States. We will continue to offer fluorescent antibody (FA) testing for Campylobacter fetus subsp. fetus, however the culture has been discontinued and replaced with Campylobacter jejuni culture. | 25.00 /sample | Bacteriology | M-S | 3-5 days | 1) Fresh feces or intestinal contents; 2) ligated loop of intestine; 3) Placenta (preferred) or fetal liver for small ruminant and camelid abortion cases; 4) Swab of 1 or 2 | 1-3) Sealed sterile separate containers (sterile preferred); 4) Swab in Amies transport media (w/charcoal preferred) | 1-2) Refrigerate |
| Canine Parvovirus Genotyping Test for genetic changes in virus. Test defines the genetic type (2a, 2b, or 2c) of canine parvovirus in clinical cases. Analysis is usually done on canine parvovirus isolates obtained from virus isolation (additional test fee). | 80.00 | Virology | As needed | 4-10 days | 1) Virus isolate; 2) Tissue sample; 3) Fecal sample | 1-3) Leakproof container | Ship in insulated container with ice pack |
| Cantharidin (Gas Chromatography/Mass Spectrometry) | 165.50 /sample | Toxicology | M-F | 1-3 days | 1) urine; 2) stomach contents, vomitus | 1-2) sterile, leak-proof container | Refrigerate w/ cool pack |
| Cerebrospinal Fluid Cytology Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation Specify if the sample is from the atlanto-occipital or lumbosacral region as result interpretation is dependent on site. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | 58.00 | Clinical Pathology | M - F | 1 business day | 1) CSF in EDTA; 2) For best results also submit freshly made, rapidly air-dried, un-stained smears. 3) Also submit a red top tube if culture anticipated | 1) Lavender-top tube; 2) Also submit a red top tube if culture anticipated | 1,3) Refrigerate fluid samples; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment. |
| Chlamydia (Chlamydia) FA Fluorescent antibody | 25.00 /animal | Virology | M-F | 2-4 days | 1) Tissue: lung; conjunctival tissue or other suspect affected tissue; fetal tissue such as placenta (preferred), liver or spleen 2) Air-dried conjunctival smear; 3) Milk | 1) leakproof, unbreakable plastic container or bag; 2) Slide mailer; 3) Leakproof container | 1,3) Ship for next day delivery in insulated container with freezer pack(s) 2) No coolant |
| Chlamydia CF Complement Fixation Species tested: bovine, ovine, caprine and feline Note: This test has not been validated on feline species. Contact laboratory before submitting samples from any other species. Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. | 11.00 /sample | Virology | Once per week | 5 days | 1 ml serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Chlamydia psittaci PCR PCR Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) culture (solid or liquid) or isolate; 2) tissue (min. 5 g); 3) feces (min. 5 g); 4) swab | unbreakable, leak-proof container | Refrigerate |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|--------------------------------|--------------------|-----------|----------|--|---|--|
| Chloride | 9.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| See Eclinpath for more information | | | | | | | |
| Chloride, urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Cholesterol | 13.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Dogs and cats should be fasted for around 12 hours to avoid cholesterol interference from post-prandial hyperlipidemia. See Eclinpath for more information | | | | | | | |
| Cholinesterase | 8.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Measurement of cholinesterase activity in serum or plasma is a quick screening test indicated for animals with a history of possible exposure to organophosphate or carbamate compounds and/or that show clinical signs compatible with exposure. Remember that erythrocytes are rich in cholinesterase, therefore hemolysis invalidates the results. | | | | | | | |
| Chromatography Quantitation | 165.50 per hour | Toxicology | M-F | 7 days | 1) 10 mL EDTA; 2) 10 mL heparin whole blood; 3) 50 mL urine | Heparinized or EDTA tube, no gel clot; sterile, leak-proof container | 1-3) Refrigerate w/ cool pack |
| Special request only, contact Lab at 607-253-3975 or -3974 | | | | | | | |
| Clostridium chauvoei FA | 14.00 /sample | Bacteriology | M-F | 1-2 days | 1) 2-inch cube of affected muscle; 2) Swab from affected lesions | 1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM) | 1) Refrigerate; 2) DO NOT refrigerate |
| For the detection of Clostridium chauvoei by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing . | | | | | | | |
| Clostridium difficile Toxins (A/B) | 33.00 /sample | Bacteriology | M, W, F | 2 days | Fresh feces or tied-off loop of bowel (with contents) | Sealed sterile container. Swabs are NOT acceptable. | Refrigerate. Fresh samples must reach the lab within 24 hours of collection. If this is not possible, freeze immediately and ship overnight frozen on dry ice. |
| For the detection of Clostridium difficile toxins A and B by ELISA in enteric samples. *May qualify for New York State contract pricing. For investigational purposes only. The sensitivity of this Clostridium difficile toxin test is maximized by freezing fecal samples that can not be sent chilled to the lab to arrive the next day, since these toxins are heat labile. If you are requesting other testing on feces or colon contents, such as parasitology exams for which the sample should not be frozen, it requires having two separate fecal specimens. The specimen held over a night or weekend can be frozen, labeled as "held frozen" and shipped with extra insulation/freezer packs. A negative toxin test result on a sample that has not been handled in this way should be interpreted with caution as it may not reflect the actual disease status of the animal. Please consider the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions. | | | | | | | |
| Clostridium novyii FA | 14.00 /sample | Bacteriology | M-F | 1-2 days | 1) 2-inch cube of affected muscle; 2) Swab from affected lesions | 1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM) | 1) Refrigerate; 2) DO NOT refrigerate |
| For the detection of Clostridium novyii by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing . | | | | | | | |
| Clostridium perfringens Enterotoxin | 28.00 /sample | Bacteriology | M, W, F | 2 days | Fresh feces or tied-off loop of bowel (with contents) | Sealed sterile container. Swabs are NOT acceptable. | Refrigerate. Fresh samples must reach the lab within 24 hours of collection. If this is not possible, freeze immediately and ship overnight frozen on dry ice. |
| For the detection of Clostridium perfringens enterotoxin by ELISA. This ELISA tests for only one of several possible toxins. Please also consider anaerobic culture. The sensitivity of this Clostridium perfringens toxin test is maximized by freezing fecal samples that can not be sent chilled to the lab to arrive the next day, since these toxins are heat labile. If you are requesting other testing on feces or colon contents, such as parasitology exams for which the sample should not be frozen, it requires having two separate fecal specimens. The specimen held over a night or weekend can be frozen, labeled as "held frozen" and shipped with extra insulation/freezer packs. A negative toxin test result on a sample that has not been handled in this way should be interpreted with caution as it may not reflect the actual disease status of the animal. Please consider the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions. | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---------------|-------------------------|-------------------|------------|---|---|--|
| Clostridium septicum FA For the detection of Clostridium septicum by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing . | 14.00 /sample | Bacteriology | M-F | 1-2 days | 1) 2-inch cube of affected muscle; 2) Swab from affected lesions | 1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM) | 1) Refrigerate; 2) DO NOT refrigerate |
| Clostridium sordellii FA For the detection of Clostridium sordelli by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing . | 14.00 /sample | Bacteriology | M-F | 1-2 days | 1) 2-inch cube of affected muscle; 2) Swab from affected lesions | 1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM) | 1) Refrigerate; 2) DO NOT refrigerate |
| Clotting Factor (05) FV Activity | 34.00 | Comparative Coagulation | M - F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (07) FVII Activity | 15.00 | Comparative Coagulation | M - F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (08) FVIII Activity Factor VIII coagulant activity | 15.00 | Comparative Coagulation | Tu-F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (09) FIX Activity Measurement of Factor IX coagulant activity in dogs and cats | 15.00 | Comparative Coagulation | Tu-F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (10) FX Activity Factor X coagulant activity in cats and dogs | 15.00 | Comparative Coagulation | M-F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (11) FXI Activity The measurement of coagulation Factor XI in cats and dogs. | 15.00 | Comparative Coagulation | Tu-F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate |
| Clotting Factor (12) FXII Activity The measurement of coagulation Factor XII in dogs and cats. | 15.00 | Comparative Coagulation | Tu-F | 1 day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigeration |
| Clotting Factor (13) FXIII Deficiency Screening Requires submission of paired sample from same-species control animal | 56.00 | Comparative Coagulation | Monday - Thursday | 1 - 3 days | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigeration |
| Coagulation Panel + Fibrinogen Incl. aPTT, PT, TCT, Fibrinogen (for small animals only) Ship Overnight: for additional info on Submission Guidelines, or see Comp Coag web page | 34.00 | Comparative Coagulation | M-F | same day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Coagulation Panel, Small Animal Incl. aPTT, PT, TCT Ship Overnight: for additional info on Submission Guidelines, or see Comp Coag web page | 27.25 | Comparative Coagulation | M-F | same day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Copper, serum/plasma Contact Lab for specifics for avian and very small animals | 25.00 /sample | Toxicology | M-F | 3 days | 1) 2 mL serum; 2) or transfer 2 mL heparinized plasma into a red top tube | 1-2) Red top tube, no gel clot | 1-2) Refrigerate. Ship in insulated container with ice pack. |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|------------------------------|-------------------------|---------------|----------|--|--|---|
| Coronavirus PCR, Alpha The alpha coronavirus PCR employed by the AHDC will detect feline enteric coronavirus, canine enteric coronavirus, ferret or mink coronavirus, and swine coronavirus - AKA transmissible gastroenteritis virus. For species specific testing information please see Coronavirus PCR Guidelines. | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) EDTA blood, serum (min. 2 mL); 2) tissue (min. 5 g); 3) ascites or body fluid (min. 2 mL); 4) feces (min 5 g) | unbreakable, leak-proof container | Refrigerate. Ship in insulated container with ice pack. |
| Corticosteroids, Drug Screen capillary GC/MS Special request only ; contact Lab at 607-253-3975 or -3974. The list of drugs covered in this screen include: Dexamethasone, Betamethasone, Methyl Prednisolone, Prednisolone, Prednisone, Flumethasone, Triamcinolone, and Triamcinolone Acetonide. | 331.00 /sample | Toxicology | M-F | 3 days | 1) 20 mL heparin whole blood; 2) or transfer 5 mL heparin plasma into a red top tube; 3) 50 mL urine | 1) Heparinized tube, no gel clot; 2) red top blood collection tube; 3) sterile container | 1-3) Refrigerate w/ cool pack |
| Cortisol baseline For more information, see Appendix C | 16.00 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate |
| Cortisol Post ACTH Post treatment sample | 10.50 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate. Ship in insulated container with ice pack. |
| Cortisol Post Dexamethasone For more information, see Endocrinology Submission Guidelines | 10.50 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate |
| Cortisol: ACTH response Pre-treatment sample For more information, see Canine Adrenal Function Tests, Equine Cushings Tests , Feline Adrenal Function Tests, or Appendix C. | 14.50 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate. Ship in insulated container with ice pack. |
| Cortisol: Dex suppression For more information, see Endocrinology Submission Guidelines | 14.50 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate |
| Creatine kinase CK, CPK See Eclinpath for more information | 8.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Creatinine See Eclinpath for more information | 11.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Creatinine, urine | 19.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs |
| Crossmatch Donor 1 Do not freeze | 42.00 per donor | Clinical Pathology | M-F | same day | 1) 1 to 3 mL EDTA whole blood AND 2) 2 mL separated serum (from each blood donor AND recipient) | 1) lavender top tube AND 2) plastic tube | Refrigerate |
| Crossmatch, Feline Ship Overnight; no NOT freeze whole blood samples | 34.75 | Comparative Coagulation | M-F | same day | 1) 2 mL EDTA whole blood AND 2) 2 mL separated serum (from each blood donor AND recipient) | 1) lavender top tube AND 2) serum tube | Refrigerate/ship on cold pack |
| Cryptosporidium ELISA ELISA plus Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate. Fecal Qualitative or Quantitative exam will be performed with each ELISA at an additional fee. This ELISA cannot be requested alone. If both Cryptosporidium and Giardia ELISA's are ordered, only one flotation will be performed and charged. | 25.00 | Parasitology | Mon, Thur | 1-3 days | feces | wide mouth leak-proof container | Refrigerate |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|------------------------------|-------------------------|---------------|----------------|---|--|--|
| Cryptosporidium Spp real-time PCR | 36.75 | Molecular Diagnostics | M-F | 3-5 days | Feces, intestinal tissue | Sealed vial or container in a secondary container | Refrigerate. Ship in insulated container with ice pack. |
| This real-time PCR assay detects the 18S rRNA gene of pathogenic Cryptosporidium species according to the method published by Operario et al. Am J Trop Med Hyg. 2015. This test is intended primarily for research applications at this time. Please contact lab for volume discounts. Ct values at or below 28 are more likely to be associated with active diarrhea. Ct values above 32 are less likely to be associated with diarrhea. | | | | | | | |
| Cyanide | 55.25 /sample | Toxicology | M-F | 2 days | 1) 10 mL EDTA whole blood; 2) 10 mL heparinized whole blood; 3) 50g stomach contents; 4) 100g forage, silage, or other material | Heparinized or EDTA tube; air-tight container | 1-2) Refrigerate; 3-4) freeze |
| 1-2) Refrigerate promptly; Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes) 3-4) freeze samples promptly, keep samples cold in transit; special request only, contact Lab to make arrangements | | | | | | | |
| Cytology Smear Exam | 48.00 | Clinical Pathology | M - F | 1 business day | 1) Smears (at least 2-3 unstained smears preferred); and/or 2) Fluid in EDTA (no cell counts or total protein provided) | 1) Frosted-end slides; 2) Lavender-top tube (if submitting fluid) | Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears/slides at room temperature; protect from chilling/coolant by insulating separately during shipment |
| Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation With fluid aspirates, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | | | | | | | |
| D-dimer Quantitative | 30.50 | Comparative Coagulation | M-F | same day | separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Ship Overnight. Rodent assay fee is \$59.50. For additional info see Comp Coag web page | | | | | | | |
| Dermatology Biopsy Surcharge | 15.00 | Anatomic Pathology | M - F | 4 - 10 days | 0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio. | Wide-mouth, leakproof, unbreakable jar; NOT GLASS | room temperature; protect from freezing |
| For Dermatology biopsies (up to 5 sites) | | | | | | | |
| DIC Panel, Small Animal | 88.50 | Comparative Coagulation | M-F | same day | 2 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Includes aPTT, PT, Fibrinogen, Antithrombin, D-dimer Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | | | | | | | |
| Dilute Russell's Viper Venom Time | 15.50 | Comparative Coagulation | M-F | same day | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | | | | | | | |
| Direct Coombs | 37.00 | Clinical Pathology | M-Sa | 1-2 days | 2-5 mL EDTA blood AND 2 dry (unfixed, unstained) blood smears | lavender top tube AND frosted-end slides | Refrigerate |
| Dog, cat, horse; coolant not in contact with specimen or slides; ship OvernightFor instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear | | | | | | | |
| Drug Quantitation | 165.50 per hour | Toxicology | M-F | 5 days | 1) 10 mL heparin plasma transferred to red top; 2) 20 mL heparin whole blood; 3) 50 mL urine | 1) red top blood collection tube; 2) Heparinized tube, no gel clot; 3) sterile container | 1-3) Refrigerate w/ cool pack |
| Special request only, contact Lab | | | | | | | |
| Ehrlichia canis PCR | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) EDTA blood, buffy coat (min. 2 mL); 2) tick: see comment; 3) tissue (min. 5 g) | unbreakable, leak-proof container | Refrigerate |
| E. canis PCR The sample required for testing from live animals is EDTA whole blood. Ticks must first be submitted for parasite ID to determine if they are a vector species; see Parasite Identification test. | | | | | | | |
| Electrolytes Panel | 17.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs |
| Na, K, Cl For more information, see Chemistry Panel Information | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|----------------------------|--------------------|-----------|-----------|--|--|--|
| Electrolytes Panel, urine Na, K, Cl, Creat For more information, see Chemistry Panel Information. | 31.00 | Clinical Pathology | M-Sa | 1-2 days | 2 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs |
| Electrophoresis, blood | 60.00 | Clinical Pathology | 1x/wk | 7-10 days | 1) 1 mL separated serum; 2) heparinized plasma for exotics/avian only | plain glass or plastic tube | Refrigerate |
| Electrophoresis, fluid All species; please specify site of fluid collection | 60.00 | Clinical Pathology | 1x/wk | 7-10 days | Body fluids. Please specify site. | plain glass or plastic tube | Refrigerate |
| Electrophoresis, urine | 60.00 | Clinical Pathology | 1x/wk | 7-10 days |) 5-7 mL urine | plain glass or plastic tube | Refrigerate |
| Enteric Bacteriology Cultures, Panel 1 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], and Salmonella culture. *May qualify for New York State contract pricing. Aerobic culture not tested on calves over 10 days of age. Antimicrobial Susceptibility (MIC) are not included in the testing fee. Please consider the Foal Diarrhea/Septicemia Kit for foal submissions. | 62.00 /sample | Bacteriology | M-Sa | 3-7 days | 1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) Swab of 1-3 | 1) Sealed, sterile container or ParaPak; 2-3) Sealed sterile container; 4) Swab in Amies transport media | 1-3) Refrigerate. Ship in insulated container with ice pack. EXCEPT ParaPak, which is shipped at room temperature and protected from temperature extremes. |
| Enteric Bacteriology Cultures, Panel 2 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], Salmonella culture and Campylobacter jejuni culture. Antimicrobial Susceptibility (MIC) are not included in the testing fee. Susceptibilities not applicable to Campylobacter jejuni. | 87.00 /sample | Bacteriology | M-Sa | 3-7 days | 1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) Swab of 1-3 | 1) Sealed, sterile container or ParaPak; 2-3) Sealed sterile container; 4) Swab in Amies transport media | 1-3) Refrigerate. Ship in insulated container with ice pack. EXCEPT ParaPak, which is shipped at room temp and protected from temperature extremes. |
| Enteric Bacteriology Cultures, Panel 3 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], Salmonella culture, Campylobacter jejuni culture, anaerobic culture [Clostridium perfringens (all species) and Clostridium difficile* (foals and primates only)] Antimicrobial Susceptibility (MIC) are not included in the testing fee. Susceptibilities are not applicable to Campylobacter jejuni and Clostridia species. *Culture for Clostridium difficile in other species available by request only. **Anaerobic Transport medias (e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM) are suitable for most aerobic cultures, including for the aerobes of concern in enteric infections. Despite labelling to the contrary, culture systems containing Amies transport media or Stuart transport media are inappropriate as transport systems for anaerobic culturing. Swabs transported without any transport media are unsuitable for both aerobic and anaerobic culture. | 132.00 /test | Bacteriology | M-Sa | 3-7 days | 1) Anaerobic transport media inoculated with swab of fresh feces or intestinal contents; or 2) Ligated ileum or jejunum; and/or 3) Mesenteric lymph node | 1) Anaerobic transport media** (BBL™ Port-a-cul™ or Anaerobic Systems ATM); or 2-3) Sealed sterile container | 1) DO NOT Refrigerate anaerobic transport media tubes or vials; 2-3) Tissues are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection. |
| Environmental Samples (for parasites) Please contact Lab: fee depends on parasite of interest, type of sample submitted, and processing required | 25.00 See comments for fee | Parasitology | M-F | 1-7 days | Please contact Lab before submitting sample to determine if sample is appropriate for testing. | | |
| Ethylene Glycol Special request only, contact Lab | 55.00 /sample | Toxicology | M-F | 2 days | 1) 10 mL heparin whole blood; 2) 10 mL EDTA whole blood; 3) urine; 4) food materials; 5) stomach contents | 1,2) Heparinized or EDTA tube, no gel clot 3-5) sterile, leak-proof container | 1-5) Refrigerate w/ cool pack |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 12 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---------------|-------------------------|-----------------|---------------|--|--|---|
| Fecal Qualitative A double centrifugation concentration fecal flotation with sugar and/or zinc sulfate solution used as a general test for evaluating feces for parasitic infections. The qualitative fecal flotation, using a double centrifugation concentration technique, is a general broad based test for evaluating feces for parasitic infections. It indicates the presence or absence of patent protozoan or worm infections in domestic and wild animals of all species. Samples from individual animals are preferred. Composite samples tell when there is a parasite problem, but not the animals that are particularly affected. In this laboratory, all samples are actively floated (centrifuged) using a sugar solution (1.33 specific gravity). Zinc sulfate solution (1.18 specific gravity) is used also for samples from small animals, animals under six months of age, and when delicate protozoa, such as Giardia, or worms, such as nematode larvae, are of concern. Using these flotation media, protozoan cysts and worm eggs and larvae are recovered for microscopic evaluation. When submitting a sample provide a sufficient sample to make a diagnosis possible, and include a history of the health concerns related to the animal(s) from which the sample(s) are collected. For best results, collect fresh samples, keep them refrigerated, and submit them for examination within seven days of collection. Samples that are preserved in 10% formalin or 70% alcohol can also be submitted, but the preservative may compromise the quality of the sample for parasite detection. | 25.00 | Parasitology | M-F | 1-3 days | 10g fresh fecal sample, animal 3 days of age or older | Plastic leakproof container | Freezer Packs. Ship Next Day Delivery |
| Fecal Quantitative Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate solution. Primarily used to examine feces of camelids, horses, and ruminants, but other species may be requested. Reported as the number of cysts/ oocysts/ eggs/ larvae per gram. Samples for animals under 3 days of age will not be tested. | 25.00 | Parasitology | M-F | 1-3 days | 10g fresh fecal sample, animal 3 days of age or older | wide mouth leak-proof container | Refrigerate |
| Fecal Wet Mount Was previously called "Direct Fecal Smear" Exam for motile parasites. This test for Cornell CUHA only (must be read within 1 hr of fecal production by host). See Trich. Fecal Exam, Staining for Other Protozoa, or Giardia ELISA | 5.50 | Parasitology | M-F | 1 day | 10g fresh fecal sample | wide mouth leak-proof container | Do NOT Refrigerate |
| Feline Blood Typing Feline blood typing Ship Overnight; do NOT freeze whole blood samples | 29.75 | Comparative Coagulation | M-F | same day | 1 to 3 mL EDTA whole blood | lavender top tube | Refrigerate/ship on cold pack |
| Feline Calicivirus (FCV) Isolation Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline. | 70.00 /animal | Virology | M-F | 21-30 days, s | 1) Nasal swab; pharyngeal swab; ocular swab; 2) Tracheal wash; broncho-alveolar lavage; 3) Upper airway tissue; lung | 1) Swabs in Viral Transport Medium; 2) Leakproof container; 3) Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Calicivirus (FCV) SN Serum neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation. | 16.50 /sample | Virology | M | 3-10 days | 1 mL serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Calicivirus FA (FCV) Fluorescent antibody | 25.00 | Virology | M - F | 2 - 4 days | | Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Coronavirus ELISA (FCOV) Feline Coronavirus ELISA (FCOV) detects antibodies to Coronavirus IgG in feline plasma or serum | 22.00 | Serology | 2-3 days/week | 2-3 days | 1ml feline serum or plasma | red top blood collection tube | Refrigerate |
| Feline Enteric Coronavirus FA Fluorescent antibody | 25.00 /animal | Virology | M-F | 2-4 days | Tissues with defined lesions | Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Enteric Coronavirus/FIP PCR The alpha coronavirus PCR employed by the AHDC will detect feline enteric coronavirus, canine enteric coronavirus, ferret or mink coronavirus, and swine coronavirus - AKA transmissible gastroenteritis virus. The coronavirus associated with Feline Infectious Peritonitis is not distinguished from enteric coronavirus of cats. FIP cases would be expected to have virus in affected tissues and fluids. For species specific testing information please see Coronavirus PCR Guidelines. Preferred sample is feces swab in red top tube or intestinal tissue. | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) EDTA blood or serum (min. 2 mL); 2) tissue (min. 5 g); 3) ascites or body fluid (min. 2 mL); 4) feces (min 5 g) Min Sample Volume: | unbreakable, leak-proof container | Refrigerate |
| Feline Heartworm Antibody Lateral Flow Immunoassay This test detects antibodies resulting from heartworm infections in cats that may include immature worms, adult male and female worms or single sex male or female infections. This test is validated for samples from domestic feline hosts only. An Occult Heartworm test that detects antigen produced by adult D. immitis can be performed at the same time using the sample submitted for the Antibody test at an additional charge. | 15.00 | Parasitology | Tuesday, Friday | 1-3 days | 0.5 mL Serum | Red top blood collection tube | Refrigerate |
| Feline Herpesvirus (FHV) (Rhinothrachetis) Isolation Virus isolation Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline. | 70.00 /animal | Virology | M-F | 21-30 days, s | 1) Nasal swab; pharyngeal swab; ocular swab; 2) Tracheal wash; broncho-alveolar lavage; 3) Upper airway tissue; lung | 1) Swabs in Viral Transport Medium; 2) Leakproof container; 3) Sealable plastic bag or container | Ship in insulated container with ice pack |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 13 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---------------|-----------------------|----------------|---------------|---|--|---|
| Feline Herpesvirus (FHV) (Rhinothracheitis) SN Serum Neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation. | 16.50 /sample | Virology | M | 3-10 days | 1 mL serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Herpesvirus (FHV) FA Fluorescent antibody | 25.00 /animal | Virology | M-F | 2-4 days | 1) lung; 2) air-dried ocular smear; 3) Fetal lung, liver, kidney, adrenal, spleen | 1) Sealable plastic bag or container; 2) Slide mailer; 3) Sealable plastic bag or container | 1, 3) ship in insulated container with ice pack 2) No coolant |
| Feline Immunodef Virus (FIV) ELISA The FIV ELISA detects antibodies to the Feline Immunodeficiency Virus (FIV). Felines with equivocal FIV ELISA results, except kittens, require Western Blot confirmation, as do all exotics. In these cases, FIVWB is added automatically unless the submitter specifies on the submission form not to run it. Note: The FIVWB cannot be ordered by itself, it must be accompanied with the FIV ELISA. If the FIVWB is requested, the FIV ELISA will be automatically added with its fee billed in addition to the FIVWB fee. | 26.00 /animal | Serology | 2x/wk | 2-3 days | 1 mL serum or plasma | red top or purple top blood collection tube | Refrigerate |
| Feline Immunodeficiency Virus (FIV) western blot The FIV Western blot detects the presence of antibodies to FIV in serum or plasma, primarily in samples having equivocal ELISA results. The FIV Western blot cannot be ordered by itself, see comment above. Felines with equivocal FIV ELISA results, except kittens, require Western Blot confirmation, as do all exotics regardless of ELISA result. In these cases, FIVWB is added automatically unless the submitter specifies on the submission form not to run it. Note: The FIVWB cannot be ordered by itself, it must be accompanied with the FIV ELISA. If the FIVWB is requested, the FIV ELISA will be automatically added with its fee billed in addition to the FIVWB fee. | 38.00 /sample | Serology | 2x/wk | 4-5 days | 2 mL serum | red top blood collection tube | Refrigerate |
| Feline Influenza Virus Isolation Virus isolation Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline. | 70.00 /animal | Virology | M-F | 21-30 days, s | 1) Nasal swab, pharyngeal swab; 2) Tracheal wash, broncho-alveolar lavage; 3) Upper airway tissue, lung | 1) Swabs in Viral Transport Medium; 2) Leakproof container; 3) Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Influenza Virus Matrix PCR This PCR assay will detect most if not all influenza virus types currently circulating in most species, including 2009 H1N1. This assay does not specifically identify the influenza H and N types. That testing can potentially be requested on positive samples if unusual outbreaks or cross-species transmission is suspected. Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. This service is performed pursuant to an agreement with Roche Molecular Systems. | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) nasal swab; or 2) tissue (min. 5 g); or 3) allantoic fluid (min. 2 mL) or other culture | unbreakable, leak-proof container | Refrigerate |
| Feline Leukemia Virus (FeLV) ELISA The FELV ELISA detects FeLV group-specific antigens in the blood of infected cats. | 19.00 /sample | Serology | M-F | 2-3 days | Serum, plasma or whole blood | red top or purple top blood collection tubes | Refrigerate |
| Feline Parvovirus (Panleukopenia) FA Fluorescent antibody | 25.00 /animal | Virology | M-F | 2-4 days | Ileum, jejunum | Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Parvovirus (Panleukopenia) Virus HI Hemagglutination Inhibition Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation. | 16.50 /sample | Virology | Varies | 2-4 days | 1 mL serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Respiratory Disease Virus Isolation Includes: FHV, FCV, FePnV, FSV + Unknowns Swabs in bacterial transport medium (Ames, Port-a-cul, etc) are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline. | 70.00 /animal | Virology | M-F | 21-30 days, s | 1) Nasal swab, pharyngeal swab; 2) Tracheal wash, broncho-alveolar lavage; 3) Upper airway tissue, lung | 1) Swabs in Viral Transport Medium; 2) Leakproof container; 3) Sealable plastic bag or container | Ship in insulated container with ice pack |
| Feline Respiratory Panel PCR Multiple assays to detect respiratory pathogens in cats Testing for feline herpesvirus and feline calicivirus can be done by virus isolation. Please request VI in addition to the Feline Respiratory Panel PCR. | 50.00 | Molecular Diagnostics | M-F, as needed | 3-5 business | nasal and oral-pharyngeal swab, lung | Securely closed tube with viral transport media or a few drops of saline | Refrigerate. Ship to lab over night in insulated container with ice pack(s) |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|----------------|-------------------------|-------------|------------|---|--|---|
| Feline Syncytial Virus (FSV) SN Serum neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation. | 16.50 | Virology | contact lab | 4-11 days | 1 mL serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Toxoplasmosis ELISA To detect Feline Toxoplasma IgG antibodies in feline serum and plasma | 20.00 | Serology | 2-3X/week | 3 days | 2 ml serum or plasma | Red top or purple top tube | Refrigerate |
| Feline Vaccine Panel 1 Includes Feline Herpesvirus (Rhinotr) SN and Feline Panleukopenia Virus HI Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. | 33.00 /sample | Virology | Varies | 5 - 9 days | 1 ml serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Vaccine Panel 2 Includes Feline Herpesvirus (Rhinotr) SN and Feline Panleukopenia Virus HI and Feline Calicivirus SN Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. | 49.50 /sample | Virology | Varies | 5 - 9 days | 1 ml serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Feline Vaccine Panel 3 Includes Panleukopenia HI and Calicivirus SN Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. | 33.00 /sample | Virology | Varies | 5 - 9 days | 1 ml serum | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| Fibrinogen Quantitative measure of clottable fibrinogen (Claus method) Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 22.00 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Fibrinogen (antigen) Fibrinogen concentration; ELISA method Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 43.00 | Comparative Coagulation | Tu, F | 1-3 days | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Field Necropsy Surcharge This charge will be added to any histopathology request on ante-mortem tissues unless deemed unnecessary by the reviewing pathologist, in which case only a standard histopathology charge will be applied. For necropsied animals it is always recommended to submit a complete set of tissues representing major organ systems and any sites potentially involved based upon clinical history, ante-mortem testing, or post-mortem gross lesions. Tissues will be selected for histopathological examination based upon the clinical history and gross description provided, at the discretion of the pathologist. A clinical history for the individual animal and potentially also for the herd or population from which the animal originates is very important. Also indicate if the animal was euthanized. | 56.00 | Anatomic Pathology | M-F | 4-10 days | 0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio. | Wide-mouth, leakproof, unbreakable jar; NOT GLASS | room temperature; protect from freezing |
| Fluphenazine (LC/MS) (Liquid Chromatography/Mass Spectrometry) Liquid Chromatography/Mass Spectrometry (LC/MS) | 165.50 /sample | Toxicology | M-F | 1-3 days | 1) 10 mL heparine whole blood; or 2) 5 mL heparine plasma transferred to red top tube | Heparinized tube, no gel clot | Refrigerate with cold pack |
| Folate, Folic Acid Avoid hemolysis and excessive exposure to direct light; see Appendix C | 27.00 | Endocrinology | Tu | 1-8 days | 1 mL serum; avoid EDTA plasma | plastic or glass tube | Refrigerate |
| Free-T4 by dialysis & TSH combination Includes Free T4 by dialysis and TSH For more information, see Appendix C | 49.00 | Endocrinology | M, Th | 1-5 days | 1 - 2 mL serum | plastic or glass tube | Refrigerate |
| Free-T4 by dialysis baseline (FT4d) For more information, see Appendix C | 42.00 | Endocrinology | M, Th | 1-5 days | 1.5 mL serum | plastic or glass tube | Refrigerate |
| Fructosamine See Eclnpath for more information | 36.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum | red top blood collection tube | Refrigerate. Ship overnight in insulated container with freezer packs |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-----------------------------|--------------------|-----------|----------------|---|---|--|
| Fungal Culture | 47.00 /sample | Bacteriology | M-F | 28+ days | 1)Nails, skin scrapings or 10-12 hairs from border of lesion; combings from asymptomatic cats (dermatophytes); 2)Fresh tissue or body fluids (systemic fungi/yeast); 3)Ear (yeast); 4) Swab of lesion, tissue, fluid or ear | 1) DRY sterile sealed container (e.g. closed urine cup or envelope); 2) Fluid or tissue in Sterile, sealed container (e.g. red top blood collection tube); 4) Swab in Amies transport media** | 1) Not required; 2) Refrigerate |
| For the isolation of yeasts, dermatophytes and other filamentous fungi. *May qualify for New York State contract pricing. **Swabs are not encouraged for tissues or body fluids. Keep tissues moist. Ship overnight. Antifungal susceptibilities are not included in the testing fee. | | | | | | | |
| Fungal ID | 47.00 /isolate | Bacteriology | M-F | 28+ days | Fungal isolate (pure culture) | Agar slant or plate (sealed) | Refrigerate |
| For the identification of pure fungal isolates (filamentous fungi or yeast). Submissions accepted from Laboratories only. Ensure purity before shipment. Follow all biohazard shipping regulations. Provide animal history and suspected identification if possible. Antifungal susceptibilities are not included in the testing fee. | | | | | | | |
| Fungal KOH Fluorescent Stain | 12.00 /sample | Bacteriology | M-F | 1-2 days | 1)Nails, skin scrapings or 10-12 hairs from border of lesion; combings from asymptomatic cats (dermatophytes); 2)Fresh tissue or body fluids (systemic fungi/yeast); 3)Prepared slide | 1) DRY sterile sealed container (e.g. closed urine cup or envelope); 2) Sterile sealed container (e.g. red top blood collection tube); 3) Slide mailer | 1,3) Not Required; 2) Refrigerate |
| For the direct microscopic examination for yeast and fungi. *May qualify for New York State contract pricing . Swabs are not acceptable | | | | | | | |
| Fungal Serology Panel | 64.00 | Bacteriology | M, W, F | 3-5 days | 1) Serum; 2) Cerebrospinal fluid (for Cryptococcus antigen only) | 1) Red top blood collection tube; 2) Sealed sterile container | 1-2) Refrigerate |
| For the detection of antibody to Histoplasma, Blastomyces, Coccidioides and Aspergillus (AGID) and Cryptococcus antigen in serum. Tests can be requested individually. See Histoplasma AGID, Blastomyces AGID, Coccidioides AGID, Aspergillus AGID or Cryptococcus antigen. | | | | | | | |
| Fungal Serology: Aspergillus Antibody | 16.00 /sample | Bacteriology | M, W, F | 3-5 days | Serum | Red top blood collection tube | Refrigerate. Ship in insulated container with ice pack. |
| For the detection of Aspergillus antibody in serum by AGID. Test is also part of the Fungal Serology Panel. | | | | | | | |
| Fungal Serology: Blastomyces Antibody | 16.00 /sample | Bacteriology | M, W, F | 3-5 days | Serum | Red top blood collection tube | Refrigerate. Ship in insulated container with ice pack. |
| For the detection of Blastomyces antibody in serum by AGID. Test is also part of the Fungal Serology Panel. | | | | | | | |
| Fungal Serology: Coccidioides Antibody | 16.00 /sample | Bacteriology | M, W, F | 3-5 days | Serum | Red top blood collection tube | Refrigerate |
| For the detection of Coccidioides antibody in serum by AGID. Test is also part of the Fungal Serology Panel. | | | | | | | |
| Fungal Serology: Cryptococcus Antigen | 22.00 /sample | Bacteriology | M-F | 1 day | 1) Serum; 2) Cerebrospinal fluid | 1) Red top blood collection tube; 2) Sealed sterile container | 1-2) Refrigerate |
| For the detection of antigen to Cryptococcus spp. in serum. Test is also part of the Fungal Serology Panel. | | | | | | | |
| Fungal Serology: Histoplasma Antibody | 16.00 /sample | Bacteriology | M, W, F | 3-5 days | Serum | Red top blood collection tube | Refrigerate |
| For the detection of Histoplasma antibody in serum by AGID. Test is also part of the Fungal Serology Panel. | | | | | | | |
| Fungal Susceptibility | 68.00 /isolate | Bacteriology | M-TW | 6 days for pur | 1) Fresh tissue or body fluids (for filamentous fungi and yeasts to assess systemic infection); 2) Fungal isolate (pure culture) – yeast or fast growing filamentous fungi ONLY | 1) Sealed sterile container (preferred) or *Amies transport media (swab); 2) Agar slant or plate (sealed) | 1) Refrigerate; 2) Not required |
| This test includes Amphotericin B, 5-Flucytosine, Anidulafungin, Caspofungin, Micafungin, Fluconazole, Itraconazole, Posaconazole, Voriconazole *May qualify for New York State contract pricing. **Swabs are not encouraged for tissues or body fluids. Susceptibilities will NOT be performed on dermatophytes or other slow growing fungi. Fungal culture MUST be performed to recover an isolate before testing can be done. ***Lag time may be longer if isolation is required. From time of isolation: 5-6 days for yeast, 12-14 days for filamentous fungi. | | | | | | | |
| Gamma Glutamyl Transferase (GGT) | 9.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | Red top (non-anticoagulant) or green top (heparin) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| See Eclinpath for more information | | | | | | | |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 16 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|------------------------------|-------------------------|-----------|----------|---|--|---|
| Gamma Glutamyl Transferase (GGT), urine | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Giardia ELISA ELISA plus Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate. A Fecal Qualitative or Quantitative exam will be performed with each ELISA at an additional fee. This ELISA cannot be requested alone. If both Cryptosporidium and Giardia ELISA's are ordered, only one flotation will be performed and charged. | 25.00 | Parasitology | Mon, Thur | 1-3 days | feces | wide mouth leak-proof | Refrigerate |
| Glucose Glucose levels in blood and fluids Delayed serum separation or submission of whole blood (cells on serum or plasma) results in a falsely decreased (uninterpretable) glucose. Serum should also be removed promptly from serum separator tubes and placed into a new red top tube. See Eclinpath for more information | 11.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum, EDTA plasma, or heparinized plasma; 2) fluid | Red top (non-anticoagulant), green top (heparin), or purple top (EDTA) tube | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Glucose, urine | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | Red top (non-anticoagulant) tube or plastic leakproof container | Refrigerate. Ship overnight in insulated container with freezer packs |
| Glutamate Dehydrogenase, blood | 8.50 | Clinical Pathology | | | 1) 1 mL separated serum or heparinized plasma; 2) fluid | | |
| Gram Stain For the direct microscopic examination for bacteria. *May qualify for New York State contract pricing. Please consider using the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions. Specific bacteria cannot be definitively identified by this test. | 8.50 /sample | Bacteriology | M-F | 1-2 days | 1) Affected tissue, lesion or fluid; 2) Prepared slide | 1) Affected tissue, lesion or fluid in sealed sterile container; 2) Slide mailer | 1) Refrigerate; 2) Not required |
| Hemoglobin (HGB) Do not allow coolant to come in contact with specimen; ship Overnight | 7.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| Hemogram (CBC), Automated (Mammalian) WBC, RBC, HgB, HCT, MCV, MCH, MCHC, RDW, Plate Count, MPV, Automated Differential Cell Count Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panelsFor instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear | 18.50 | Clinical Pathology | M-Sa | 1-2 days | 1) EDTA blood (> 1/2 full) AND 2) 2 dry (unfixed, unstained) blood smears | 1) lavender top tube; 2) frosted-end slides | 1) Refrigerate blood; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment. |
| Hemogram (CBC), Partial WBC, RBC, HgB, Platelet Count, PCV, MPV, RDW, MCH, MCHC, MCV Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Hematology Testing. | 11.50 | Clinical Pathology | M-Sa | 1-2 days | 1) EDTA blood (> 1/2 full) AND 2 dry, unfixed, unstained blood smears | 1) lavender top tube; 2) frosted-end slides | Refrigerate |
| Hemophilia Panel, Small Animal Incl. aPTT, PT, TCT, Factor VIII:C, Factor IX:C, vWF:Ag. See interpretation information. Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 69.75 | Comparative Coagulation | M-F | same day | 2 - 3 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Hemostasis Panel, Small Animal Incl. aPTT, PT, TCT, vWF:Ag. See interpretation information Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 44.50 | Comparative Coagulation | M-F | same day | 2 - 3 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Heparin (Low Molecular Weight) Anti-Xa activity of low molecular weight heparin Ship Overnight; for determination of peak anti-Xa values, collect sample 2-3 hrs (cat) or 3-4 hrs (dogs) after heparin dose. For additional info on Submission Guidelines, or see Comp Coag web page. Also see Heparin Assay FAQs. | 39.25 | Comparative Coagulation | M-F | same day | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---|-------------------------|-----------|-----------|--|---|---|
| Heparin (Unfractionated) Anti-Xa activity of unfractionated heparin Ship overnight; specify UFH or LMWH; for additional info on Submission Guidelines, or see Comp Coag web page. Also see Heparin Assay FAQs. Peak and trough values: samples for determination of Peak anti-Xa activity should be collected at 3 to 4 hours post treatment (dogs) or 2 to 3 hours post treatment (cats). Samples for Trough levels should be collected just before the next scheduled dose. | 39.25 | Comparative Coagulation | M-F | same day | 0.5 to 1.0 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Histopathology Microscopic examination of formalin fixed, paraffin embedded tissue slides with hematoxylin and eosin staining. Additional special stains, including up to two immuno-histochemistry (IHC) stains, will be added if deemed necessary by the pathologist. Additional IHC stains may be requested for an additional fee. The 1-2 site fee is intended for surgical biopsies and is the baseline charge for the special biopsy categories below. Surcharges for these will be added to the surgical biopsy fee based on the following criteria: Dermatology biopsies: Up to five sites or tissue punches. Please use the Dermatohistopathology form to request this service. For more information, please see Dermatopathology Samples. Liver Biopsy: All canine and feline liver biopsies submitted through the surgical pathology service will receive a surgical biopsy charge plus an additional \$45.00 surcharge for a liver special stain panel. This additional charge will cover a series of special stains and review by our liver pathology specialist since the H&E stain alone may be insufficient to provide proper hepatic analysis. Upon request, a Digital Copper Quantification (DCQ) and/or consultation with a clinical specialist in hepatic disease can be added for additional fees. Please use the Small Animal Liver Biopsy form to request these services. Due to the need for these special stains the processing of reports for these specimens may not be available for 7-10 business days. Special/Augmented biopsies: Intended for complex submissions of entire organs, lobes, systems, tumor beds, limbs, jaws, etc. Field Necropsy: For necropsied animals it is always recommended to submit a complete set of tissues representing major organ systems and any sites potentially involved based upon clinical history, ante-mortem testing, or post-mortem gross lesions. Tissues will be selected for histopathological examination based upon the clinical history and gross description provided, at the discretion of the pathologist. A clinical history for the individual animal and potentially also for the herd or population from which the animal originates is very important. Also indicate if the animal was euthanized. Please contact the lab to arrange for STAT testing. Three tissues per case will be reviewed as a STAT unless prior arrangements are made. Call 607-253-3319 or email pathology service @cornell.edu. *May qualify for New York State contract pricing | 66.50 for 1-2 sites; \$20.50 for each additional site. Dermatology biopsies add \$15.00 surcharge (up to 5 sites), liver biopsies add \$45 surcharge, Special/Augmented or Field Necropsies add \$56.00. If STAT call Lab. | Anatomic Pathology | M-F | 4-10 days | 0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio. | Wide-mouth, leakproof, unbreakable jar; NOT GLASS | room temperature; protect from freezing |
| Histopathology (Each additional site) Additional charge added for each site evaluated beyond 2 that are part of a standard biopsy. These additional fees will be applied to submissions of multiple masses from a single organ that require separate slides to track specific anatomic locations and monitor margins. They will also apply to submissions for research and legal purposes, as well as general purposes, for which every submitted tissue must be examined. Please call the lab to arrange STAT testing. Three tissues per case as a STAT unless prior arrangements are made. Please call 607-253-3319 to arrange STAT histopathology. | 20.50 each additional site | Anatomic Pathology | M-F | 4-10 days | 0.5 - 1 cm thick section of tissue fixed in 10% neutral buffered formalin, 10:1 formalin to tissue ratio | Wide-mouth, unbreakable jar containing 10% neutral buffered formalin; NOT GLASS | |
| Histopathology, Liver Special Stain Panel Liver Special Stain fee is in addition to the Histopathology fee. \$100 Stat fee will be charged for nights, weekends or holidays. Call 607-253-3319 to arrange. Stat fees waived for zoonotic disease suspects or herd health outbreaks. Liver special stains will be added to every canine/feline liver biopsy case submitted for histopathology. These liver cases are reviewed by a liver pathology specialist, and consultation with a clinical specialist in hepatic disease. | 45.00 per block | Anatomic Pathology | M-F | 7-10 days | 2 cm thick section of liver tissue fixed in 10% formalin (10 times volume of sample) | Unbreakable, wide-mouth jar containing formalin; NOT GLASS | |
| Histopathology, Slide Review (1-2 slides) 1-2 slides Contact Path Dept (607-253-3312) | 55.00 | Anatomic Pathology | M-F | 3 days | 1 to 2 Slides | Slide mailer | |
| Histopathology, Slide Review (3+ slides) 3 or more slides Contact Path Dept (607-253-3312) | | Anatomic Pathology | M-F | 3 days | 3 or more slides | Slide mailer | |
| IHC w/o Histopathology | 35.00 | Anatomic Pathology | M-F | 2-7 days | formalin fixed tissue (paraffin-embedded), or tissue mounted on glass slides | | room temperature; protect from freezing |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-----------------|-------------------------|---------------|----------|--|--|--|
| Immunoglobulin G, Canine and Feline IgG by radial immunodiffusion (RID). Dog and cat only (note, bovine, horse and camelid IgG is now measured using a different method; see specific IgG measurement for those species). | 30.00 | Clinical Pathology | 1x/week | 7 days | 500 uL separated serum | plain glass or plastic tube | Refrigerate |
| Immunoglobulin G, Colostrum (Canine and Feline) IgG by radial immunodiffusion (RID). Canine and Feline only. | 36.00 | Clinical Pathology | M-Sa | same day | 1 ml | plain glass or plastic tube | Refrigerate |
| Influenza Virus Matrix PCR This PCR assay will detect most if not all influenza virus types currently circulating in most species, including 2009 H1N1 and the 2015 Canine H3N8. This assay does not specifically identify the influenza H and N types. That testing can potentially be requested on positive samples if unusual outbreaks or cross-species transmission is suspected. Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. All canine and feline samples testing positive for influenza A matrix will be further tested to determine subtype. Subtyping will be done automatically and at no additional charge. Results will be available within 3-5 days following release of the original result on an addended report. *May qualify for New York State contract pricing. This service is performed pursuant to an agreement with Roche Molecular Systems. | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) nasal swab; or 2) tissue (min. 5 g); or 3) allantoic fluid (min. 2 mL) or other culture | unbreakable, leak-proof container | Refrigerate |
| Insulin baseline - not equine Avoid hemolysis and lipemia. For more information, see Appendix C | 19.00 | Endocrinology | M, W, Th | 2-6 days | 1.5 mL serum or EDTA plasma; do not send sample on cells | plastic or glass tube | Refrigerate |
| Ionized Calcium Test run on serum only For special sample collection instructions, see Appendix A | 18.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum ONLY (see Appendix A) | non-anticoagulant tube (plain red top) | Refrigerate |
| Ionophore Quantitation Monensin, Lasalocide, Salinomycin Liquid Chromatography/Mass Spectrometry (LC/MS) | 165.50 per hour | Toxicology | M-F | 1-3 days | 1) feed; 2) 10 mL heparin whole blood | 1) leak-proof zip-lock bag; 2) green top blood collection tube | Refrigerate w/ cool pack |
| Ionophore Screen Monensin, Lasalocide, Salinomycin Liquid Chromatography/Mass Spectrometry (LC/MS) | 330.75 /sample | Toxicology | M-F | 1-3 days | feed samples | leak-proof zip-lock bag | Refrigerate w/ cool pack |
| Iron | 8.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma; | plain glass or plastic tube | Refrigerate |
| Iron Panel Iron (FE), Total Iron Binding Capacity, % Saturation for additional info on Clin Path panels | 20.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Kaolin Clotting Time (KCT) Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 15.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Knott's Heartworm Used to assess microfilarial burden or to differentiate Dirofilaria immitis and Dipetalonema reconditum microfilariae. If an Occult heartworm antigen ELISA test is also desired, the laboratory can separate plasma from the EDTA or heparin whole blood (if sufficient volume) or a serum sample can be submitted. Please clearly indicate each test as a unique test request. | 16.50 | Parasitology | M-F | 1-3 days | 1 mL EDTA or heparin whole blood | sterile blood tube | Refrigerate |
| Lead, blood Lead, blood Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes). For avian and exotics, contact Laboratory | 35.00 /sample | Toxicology | M-F | 2 days | 1) EDTA whole blood (min .5 ml); 2) heparin whole blood (min .5 ml) | Heparinized or EDTA tube | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| Lead, liquids/fluids | 30.00 /sample | Toxicology | M-F | 1-2 days | 1) urine; 2) water; 3) milk | 1-3) sterile container | 1-3) Refrigerate w/ cool pack |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 19 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-----------------------------|-----------------------|-----------|----------|---|---|--|
| Lead, tissue/feed/other (Atomic Absorption Spectroscopy - Graphite Furnace) | 45.00 | Toxicology | M-F | 3 days | 1) 2g tissue; 2) 100g feed | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |
| Leptospira MAT, 5 standard serovars | 16.00 /5 serovars | Bacteriology | M, W, F | 2-3 days | 2.0 ml serum | Red top blood collection tube | Refrigerate. Ship overnight in an insulated container with ice packs |
| For the detection of antibodies to Leptospira interrogans serovars Pomona, Hardjo, Icterohemorrhagiae/copenhageni, Grippotyphosa, and Canicola by serum microagglutination. Please indicate any previous testing at the AHDC for accurate test result interpretations. Testing paired samples at an interval between 2-6 weeks between blood draws has more diagnostic significance than relying on the results from a single sample. See also Leptospira MAT, 7 serovars for additional serovar options that may be relevant. The first 5 serovars may qualify for New York State contract pricing. Refer to Leptospira Microagglutination Testing at the AHDC for more information. | | | | | | | |
| Leptospira MAT, 7 serovars | 30.00 /7 serovars | Bacteriology | M, W, F | 2-3 days | 2.0 ml serum | Red top blood collection tube | Refrigerate. Ship overnight in an insulated container with ice packs |
| For the detection of antibodies to Leptospira interrogans serovars Pomona, Hardjo, Icterohemorrhagiae/copenhageni, Grippotyphosa, Canicola, Autumnalis and Bratislava by serum microagglutination. This test adds Leptospira interrogans serovars Bratislava and Autumnalis to the Leptospira MAT, standard 5 serovars. These two serovars are much less likely to be found in the United States, however, the AHDC has identified clinical samples with strongly elevated titers to one or both of these serovars in the absence of strong reactions to the other five. Testing for these two serovars may therefore be of diagnostic value. Please indicate any previous testing at the AHDC for accurate test result interpretations. Testing paired samples at an interval between 2-6 weeks between blood draws has more diagnostic significance than relying on the results from a single sample. Refer to [Leptospira Microagglutination Testing | | | | | | | |
| Leptospira real-time PCR | 36.75 per specimen tested | Molecular Diagnostics | M-F | 3-5 days | 1) Urine; or 2) Blood (EDTA whole blood); or 3) Urine and blood (EDTA whole blood); or 4) Kidney and/or liver; or 5) Fetal lung, kidney, liver, stomach or abomasal contents, placenta; or 6) Ocular fluids or 7) Bacterial culture. Refer to Testing Strategy information below. | Leak-proof container | Refrigerate. Ship in insulated container with freezer pack |
| This assay targets the lipL32 gene encoding the major surface antigen of pathogenic Leptospira species. | | | | | | | |
| Lipase | 24.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| Listeria Culture | 22.00 /sample | Bacteriology | M-F | 5-7 days | 1) Encephalitic: brain stem, cerebellum; 2) Visceral: liver, spleen, kidney; 3) Swab of 1 pr 2; 4) Silage | 1-2) Sealed separate sterile containers; 3) Amies transport media (swab); 4) Sealed sterile container | 1-3) Refrigerate |
| For the culture of Listeria monocytogenes ONLY. Neurologic listeriosis is not typically associated with a bacteremia. CNS infection is believed to and has been shown to result from localized entry of the bacteria via defects in the oral mucosa, with initial entry into the CNS via the trigeminal nerve. A systemic or bacteremic phase is not usually associated with this process. CSF or brain would be required to identify neuroinfection in most cases. Blood can be a good sample in monogastrics and neonates with septicemia. Consider submission of inoculated Blood Culture media and requesting "Blood culture, aerobic" for testing a clinically ill live animal. Placenta and fetal tissues are appropriate specimens in abortions cases. Our laboratory does not provide food safety testing for listeria on milk or other foods. Samples submitted for testing should be related to clinical investigations. Antimicrobial susceptibilities are not included in the testing fee. Test is also part of Ruminant CNS Panel. Rabies testing should also be submitted to the NYS DOH Rabies laboratory for neurologic submissions. Preliminary results will be provided one day after sample processing begins. | | | | | | | |
| Magnesium | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| Magnesium, urine | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| Mass Spectroscopy Confirmation | 165.50 /sample | Toxicology | M-F | 5 days | 1) 10 mL heparin whole blood; 2) 20 - 50 mL urine; 3) tissue; 4) plant materials; 5) stomach contents | 1) Heparinized tube; 2-5) sterile, clean, pesticide-free container | 1-2) Refrigerate w/ cool pack; 3-5) freeze tissue |
| Freezer pack whole blood: SPECIAL REQUEST - PLEASE CONTACT LAB. | | | | | | | |
| Mass Spectroscopy Screen | 331.00 per sample | Toxicology | M-F | 7 days | 1) 10 mL heparin whole blood; 2) 20 - 50 mL urine; 3) tissue; 4) plant materials; 5) stomach contents | 1) Heparinized tube; 2-5) sterile, clean, pesticide-free container | 1-2) Refrigerate w/ cool pack; 3-5) freeze tissue |
| Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | | | | | | | |
| Mercury | 55.00 | Toxicology | M-F | 3 days | 1) 2g tissue; 2) 100g feed; 3) 2 mL heparin whole blood | 1) sterile container; 2) any container; 3) Heparinized or EDTA tube | 1, 3) Refrigerate; 2) none |
| (Flameless Atomic Absorption) Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes). | | | | | | | |
| Microscopic Feed Analysis | 40.00 /hr | Toxicology | M-F | 3 weeks | 100g feed | any container | None |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---|--------------------|-----------|-----------|---|---|-------------------------------|
| Mineral Panel (Tissue, Feed) (Inductively Coupled Plasma Atomic Emissions Spectroscopy) Includes Ca, P, Mg, Na, K, Cu, Fe, Mn, Zn, Mo | 47.00 | Toxicology | 1x/wk | 3-7 days | 1) 2g tissue; 2) 100g feed. Do not submit serum or plasma. | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |
| Mineral Panel, urine Urine mineral panel includes Magnesium, Phosphorous, Creatinine and Calcium For additional info on Clin Path panels | 33.00 | Clinical Pathology | M-Sa | 1-2 days | 2 mL urine | plain glass or plastic tube | Refrigerate |
| Mineral/Electrolytes Panel Na, K, Cl, Bicarb, Anion Gap, Ca, P, Mg For additional info on Clin Path panels | 21.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Mineral/Electrolytes Panel, urine Na, K, Cl, Ca, P, Mg For additional info on Clin Path panels | 39.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| Mycoplasma Culture For the culture of Mycoplasma species. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Test is also part of the Reproductive Panel 4 and Respiratory Panel 6. | 27.00 /sample | Bacteriology | M-F | 5-10 days | 1) Uterine, cervical or vaginal mucus; 2) Semen; 3) Tracheal washing or lung tissue; 4) Fluid (joint, pericardial, urine, etc.); 5) Eye; 6) Ear; 7) Aseptically collected swab of 1-6 | 1-6) Tissue or fluid in sealed sterile container 7) Swab in Amies transport media WITHOUT charcoal or other suitable mycoplasma culture transport media | 1-6) Refrigerate |
| Mycotoxin Multiresidue Screen Aflatoxins (B1, B2, G1, G2), Ochratoxin A, Zearalenone, T-2 Toxin, DAS, DON and Sterigmatocystin The AHDC is no longer providing mycotoxin testing. We are recommending submission to the North Dakota State University Veterinary Diagnostic laboratory for mycotoxin testing. http://www.vdl.ndsu.edu/ | 165.50 | Toxicology | | | | | |
| Necropsy After Hours Fee An after-hours fee of \$50 will be applied when bodies are dropped off after 6pm during the week or between 3pm Saturday and 7:30am Monday morning. Additional STAT fees may apply if a necropsy is required at the time of drop off. | 50.00 | Anatomic Pathology | | | | | |
| Necropsy, 0.25 to 15 lbs Includes cats, dogs, non-contract covered poultry, rabbits, ferrets, most birds, small animal abortions, and non-contract bovine and small ruminant abortions \$100 Stat fee will be charged for nights, weekends or holidays. Stat fees will be waived for zoonotic disease suspects or herd health outbreaks. Please call 607-253-3319 to arrange. For shipping instructions, please see Shipping Necropsy Specimens. To ship by UPS or FEDEX, package must weigh less than 150 lbs. They require notification prior to scheduled pickups for all packages exceeding 75 lbs. | 125.00 in state; \$200 out of state. Additional STAT fee may apply. Contact lab for more information | Anatomic Pathology | M-F | 2-12 days | carcass | | |
| Necropsy, 16 to 40 lbs Includes small dogs, raccoons, large cats \$100 Stat fee will be charged for nights, weekends or holidays. Stat fees will be waived for zoonotic disease suspects or herd health outbreaks. Please call 607-253-3319 to arrange. For shipping instructions, please see Shipping Necropsy Specimens. To ship by UPS or FEDEX, package must weigh less than 150 lbs. They require notification prior to scheduled pickups for all packages exceeding 75 lbs. | 260.00 in state; \$350 out of state. Additional STAT fee may apply. Contact lab for more information | Anatomic Pathology | M-F | 2-12 days | carcass | | |
| Nematode Larval Culture Requires 14-day lag time from submission; Fecal Quantitative test done first to determine need | 55.00 | Parasitology | M-F | 14+ days | 15g fresh fecal sample | wide mouth leak-proof | Do not refrigerate sample |
| Nitrate/Nitrite, liquids/fluids | 24.00 /sample | Toxicology | M-F | 2 days | 1) water; 2) serum; or 3) transfer heparin or EDTA plasma to a red top tube | 1) sterile leak-proof container; 2-3) red top blood collection tube (no gel separator tubes, please) | 1-3) Refrigerate w/ cool pack |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 21 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|-------------------|--------------------|-----------|----------|--|---|--------------------------------------|
| Nitrate/Nitrite, solids | 33.00 /sample | Toxicology | M-F | 2 days | 1) stomach or rumen contents; 2) feeds | 1) sterile leak-proof container; 2) leak-proof zip-lock bag | 1-2) Freeze |
| Non-Esterified Fatty Acid (NEFA) NEFA for additional info on Clin Path panels | 13.00 | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum; 2) EDTA plasma | 1) red top blood collection tube; 2) purple top collection tube | Refrigerate |
| Occult Blood (Coprology) Do not allow coolant to come in contact with specimen; ship Overnight | 16.00 | Clinical Pathology | M-Sa | 1-3 days | feces | sterile container | Refrigerate |
| Occult Heartworm The Synbiotics DiroCHEK heartworm test is an indirect enzyme-linked immunosorbent assay (ELISA) used for the detection of adult female <i>Dirofilaria immitis</i> antigen in canine and feline serum or plasma. This test is validated for use in domestic or wild dogs and cats only. Other animals may be tested by this method, but reactivity in other species is not clearly defined. This test method can be used to detect <i>Dirofilaria immitis</i> five to six months after infection in dogs, and seven to eight months after infection in cats. Because of the nature of this disease in cats, veterinarians may want to consider including feline heartworm antibody testing for a more accurate assessment of a cat's true infection status. If a Knott's heartworm test (microfilarial detection) is also desired, EDTA or heparin whole blood (1.0 mL) is needed. Please clearly indicate each test as unique test requests. | 17.00 | Parasitology | Tu, F | 1-7 days | 1 mL serum or 1 mL EDTA or heparin plasma | sterile blood tube | Refrigerate |
| Organochlorine Insecticide Screen Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | 331.00 /sample | Toxicology | M-F | 5 days | 1) 10 - 20 mL heparin whole blood; 2) 50 mL urine; 3) 50g stomach or rumen contents; 4) 50g liver; 5) 200g other materials | 1) Heparinized tube, no gel clot; 2) sterile, leak-proof container; 3-5) sterile glass container | 1-5) Refrigerate w/ cool pack |
| Organophosphate/Carbamate Insecticide Screen Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | 331.00 per sample | Toxicology | M-F | 5 days | 1) 10 - 20 mL heparin whole blood; 2) 50 mL urine; 3) 50g stomach or rumen contents; 4) 50g liver; 5) 200g other materials | 1) Heparinized tube, no gel clot; 2) sterile, leak-proof container; 3-5) sterile glass container | 1-5) Refrigerate w/ cool pack |
| Osmolality Do not allow coolant to come in contact with specimen | 19.50 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine; 4) milk replacer | plain glass or plastic tube | Refrigerate |
| Packed Cell Volume PCV or HCT (Hematocrit) Do not allow coolant to come in contact with specimen; ship Overnight | 7.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| Parasite Identification Endo- and ecto- parasites All ticks submitted for PCR (Lyme, Anaplasma, Ehrlichia) must be identified prior to testing. A biopsy is examined directly or undergoes pepsin digestion to recover nematodes. Gastrointestinal contents and other tissues such as lung are examined and processed as needed to collect and identify parasites present. Intestinal smears are made from fresh, non-preserved intestinal sections and are used to identify non-patent coccidian infections postmortem Other samples that can be submitted for parasite identification include: Biopsies from tissues that may contain parasites; Necropsy tissues, such as sections of organs and intestines with contents; Fluids such as tracheal and cloacal washes, duodenal aspirates, urine, blood; Histological sections on slides; Free living worms or arthropods found on animals, in residences, in feed samples and environmental samples such as soil and composted material. | 28.00 | Parasitology | M-F | 1-7 days | whole parasites including worms or arthropods; punch biopsy, skin or conjunctival snips, fresh tissue for worm identification; impression smears; stained tissue section slides; environmental samples | plastic leak proof container, such as blood tube, appropriate for sample. Submit fresh, in 70% ETOH or 10% formalin, depending upon the sample. | Refrigerate if not fixed |
| Parvovirus Real Time PCR Real time PCR for the detection of parvovirus in carnivores Real time PCR for parvovirus detects presence of wild type virus as well as vaccine strains of the virus. | 36.75 | Virology | as needed | 3-5 days | 1) 5cc feces; 2) fecal swab; 3) ileum, jejunum, tongue, heart | 1) Sealable plastic bag or container; 2) Leak-proof container | Refrigerate. Ship with freezer packs |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|-----------------------------|-------------------------|---------------------------------------|----------------|--|--|--|
| Pericardial Fluid Cytology | 50.00 | Clinical Pathology | M - F | 1 business day | 1) Pericardial fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment. |
| Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With pericardial fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | | | | | | | |
| Peritoneal Fluid Cytology | 50.00 | Clinical Pathology | M - F | 1 business day | 1) Peritoneal fluid in EDTA; 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only, DO NOT refrigerate smears |
| Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With peritoneal fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | | | | | | | |
| Pesticide Screen | 331.00 /sample | Toxicology | M-F | 5 days | 1) 20 heparin whole blood; 2) 50 mL urine; 3) 500 mL water; 4) 50g liver; 5) stomach contents; 6) 100g feed; 7) plant material | 1) Heparinized tube, no gel clot; 2-7) sterile leak-proof container; | 1-3) Refrigerate w/ cool pack; 4-7) freeze |
| Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis. | | | | | | | |
| Phenobarbital | 33.00 | Clinical Pathology | M-Sa | 1 day | 1) 1 mL Serum, 2) EDTA plasma, 3) Heparin plasma | 1) Red top tube, 2) Corvac, 3) Lavender top tube, 4) Green top tube | Refrigerate, ship on ice packs |
| Phenobarbital is used for treatment for forms of epilepsy. This test measures the amount of phenobarbital in the sample Please refer to the Phenobarbital Chemistry Serum Test Expanded Information Sheet for more information. | | | | | | | |
| Phosphate (Phosphorus) | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| Phosphate (Phosphorus), urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| Platelet Aggregation | 97.00 | Comparative Coagulation | --- | --- | requires special arrangement with lab | | |
| Combination of individual tests Special arrangement - contact laboratory to schedule testing | | | | | | | |
| Platelet Count | 18.00 | Clinical Pathology | M-Sa | 1-2 days | 1) EDTA blood (> 1/2 full); 2) 2 dry (unfixed, unstained) blood smears | 1) lavender top tube; 2) frosted-end slides | Refrigerate |
| Do not allow coolant to come in contact with specimen or slides; ship Overnight | | | | | | | |
| Platelet Flow Cytometry-1 | 84.00 | Comparative Coagulation | Requires prior arrangement -once/week | 2 - 5 days | 1.5 mL EDTA blood, Citrate blood | EDTA or citrate collection tube | Do Not Freeze |
| Cytometric screen of platelet function Pre-approval by direct phone consultation with the coagulation lab is required to coordinate sample submission | | | | | | | |
| Platelet Flow Cytometry-2 | 122.00 | Comparative Coagulation | | 2 - 5 days | 3.0 mL citrate blood or platelet concentrate | On-site collection (citrate blood) | Do Not Freeze |
| Extended Platelet Cytometry Panel Platelet flow 2 includes activation and labeling reactions to define multiple parameters of platelet function, reserved for samples collected on-site (at AHDC) , or evaluation of platelet concentrates for transfusion | | | | | | | |
| Platelet Function | 52.00 | Comparative Coagulation | | 2-3 days | requires referral appointment | | |
| Combination of individual tests Special arrangement - contact laboratory to schedule testing | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|-----------------------------|-------------------------|---------------|-----------------------------------|--|--|--|
| Pleural Fluid Cytology | 50.00 | Clinical Pathology | M - F | 1 business day | 1) Pleural fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment. |
| Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With pleural fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | | | | | | | |
| Pneumovirus N RT-qPCR | 36.75 | Molecular Diagnostics | M-F as needed | 3-5 days | 1) Nasal swab, pharyngeal swab; 2) Tracheal wash, broncho-alveolar lavage; 3) Upper airway tissue, lung | 1) Swabs in Viral Transport Medium; 2) Leakproof container; 3) Sealable plastic bag or container | Freezer Packs |
| Identifies pneumovirus in dogs and cats. The virus has been known since the late 1930's but it was long believed to infect only rodents. The virus has been known since the late 1930's but it was long believed to infect only rodents. It is now commonly found in dogs and cats during outbreaks of acute respiratory disease, particularly when animals are in close confinement. There is no established etiologic role for pneumovirus in these outbreaks as multiple pathogens are typically involved. Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. This service is performed pursuant to an agreement with Roche Molecular Systems. | | | | | | | |
| Potassium | 8.50 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| Potassium, urine | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| PreAnesthesia Panel, Small Animal | 29.00 per test | Clinical Pathology | M-Sa | 1-2 days | 2 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| for additional info on Chemistry Panel Information | | | | | | | |
| Prekallikrein Screen | 14.50 | Comparative Coagulation | M-F | same day | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Ship Overnight: requires aPTT on same sample; For additional info on Submission Guidelines, or see Comp Coag web page | | | | | | | |
| Progesterone baseline | 14.50 | Endocrinology | see comment | 1-3 days | 1 - 1.5 mL serum; separated within 2 hrs after blood collection; avoid SST tubes | plastic or glass tube | Refrigerate |
| Dogs and llamas tested daily M-F; other species, M, W, F. For more information, see Endocrinology Testing and Protocols | | | | | | | |
| Protein C | 38.75 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Protein C Ship Overnight; For additional info on submitting samples see Submission Guidelines. Also see Protein C FAQs | | | | | | | |
| Prothrombin Consumption Index | 40.00 | Comparative Coagulation | Th-F | 1-2 days | paired citrate plasma and serum samples | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Ship Overnight; tested by special arrangement: contact laboratory for instructions | | | | | | | |
| Prothrombin Time (PT) | 14.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | | | | | | | |
| Rabies Brain Extraction | 51.00 | Anatomic Pathology | M-Sa | 24 hrs, follow head or whole body | | leak-proof container | Refrigerate, do not freeze |
| This procedure is intended for removal of the brain from the skull for further rabies testing and other diagnostics. A Rabies History Form is required for all rabies suspect cases. Submitting the head or carcass to the Anatomic Pathology service for brain extraction in rabies suspect cases will be followed by shipment of brain tissue to the NYS Dept of Health Lab for Rabies testing. Consideration for field collection of brain specimens and direct shipment to the NYSDOH by attending veterinarians should be considered in case involving human exposure, especially when weekends and holidays will result in further shipping delays. See these links for further information on collection and submission of appropriate specimens for direct submission. | | | | | | | |
| Red Blood Cell (RBC) Count, Total | 7.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| Do not allow coolant to come in contact with specimen; Ship Overnight. | | | | | | | |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 24 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|--------------------------------------|--------------------|---------------|---------------|--|---|---|
| Red Blood Cell (RBC), Parasites This test is performed using Wright's (Giemsa) Stain. Do not allow coolant to come in contact with specimen; ship Overnight | 12.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| Ref: Acetylcholine Receptor Antibody (Myasthenia Gravis) Immunoprecipitation radioimmunoassay for detection of circulating antibodies against nicotinic Acetylcholine Receptors (AChR) associated with canine and feline Myasthenia Gravis. This test can be performed on Canine and Feline samples only. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 120.00 (+ referral fee; see comment) | Referral | M-F | 7-9 business | 2 mL serum | Red top blood collection tube | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Adrenal Androgen Panel (Sex Hormone Panel) Canine or Fel Baseline and post-ACTH cortisol, progesterone, 17-hydroxyprogesterone, estradiol, androstenedione, testosterone This test can be performed on Canine and Feline samples only. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 180.00 (+ referral fee; see comment) | Referral | M - F | 9-12 business | 2ml each Pre and Post ACTH Serum (1 hr. post ACTH; 2 hr. if ACTH gel) – No Plasma; Allow sample to clot then centrifuge and separate serum | Red top blood collection tube (no serum separator tubes) | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Aldosterone Assay to quantitate aldosterone levels in EDTA plasma or serum. Can test a single sample for baseline values or can provide pre and post ACTH samples as ACT stimulation test. For All Species: Pre or post samples can be serum, plasma or one of each. Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 40.00 (+ referral fee; see comment) | Referral | Mon and Thurs | 4-7 Business | 1mL EDTA plasma or serum; serum or EDTA plasma should be separated from clot or cells by centrifugation, respectively, within 30 minutes, and transferred to plain tube with no additives. | Red top, Purple top, or Tiger top blood collection tube | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Allergy Testing, Can and Fel The AHDC is no longer providing referrals for this testing. The testing labs offer customizable testing which is best selected by the clinician in direct contact with the testing laboratory. Please see the referral information tabs for the laboratory to which we had previously referred testing. | 0.00 (+ referral fee; see comment) | Referral | | | | | |
| Ref: Amino Acids Complete amino acid analysis in plasma. Reference ranges available for canine, feline and equine Send via overnight courier on dry ice; Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 168.00 (+ referral fee; see comment) | Referral | W | 7-12 days | 2 mL Heparinized plasma/frozen. Separate and freeze ASAP | Insulated container, If glass tube is used, protect from breakage with secondary container or wrap. | Dry ice |
| Ref: Aspergillus Panel This panel includes Aspergillus Antibody Elisa, Electrophoresis to Quantitate host plasma proteins, and galactomannan assay for a dominant Aspergillus antigen. Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Testing performed on Avians and Mammals. Sample should be 5 days old or less at time of testing. Do not collect and ship to AHDC prior to 3 day weekend or other conditions which will not allow delivery to referral lab for testing by 5 days post collection. For test interpretations, please see Testing for Avian Aspergillus | 96.00 | Referral | M - F | | 0.5 ml heparinized plasma | Green top blood collection tube | Refrigerate, ship sample over night in insulated container with freezer packs. |
| Ref: Babesia Gibsoni Giemsa Smr slides, Exp Evaluation of blood smears for hemoparasites/ hemotropic bacteria. Also known as Hemoparasite exam – Wright Giemsa Stain Submitter must make blood smear slides. Do not submit EDTA whole blood. Testing can be performed on the following: Amphibian, Avian, Bovine, Canine, Caprine, Cervid, Equine, Exotic, Feline, Ovine, Porcine, Reptile, and Rodents. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 19.00 (+ referral fee; see comment) | Referral | Monday-Friday | 2 days | 1-2 Unstained, air-dried blood smear slides made from EDTA Whole Blood | slide mailer | Hold at room temperature. Ship at room temperature. Protect from cold temperatures with insulation. |
| Ref: Babesia PCR This is a polymerase chain reaction assay which amplifies and detects target nucleic acid sequences of the Babesia genus. This laboratory performs nucleic acid sequencing for all positive samples, verifying detection and providing identification to the species and occasionally strain level. Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. This test can be performed on all species. | 60.00 (+ referral fee; see comment) | Referral | Tues; Thurs | 5-7 days | 2ml EDTA Whole Blood | Purple Top Collection Container | Refrigerate. Ship in insulated container with freezer packs to arrive next day. |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-------------------------------------|------------|-------------------|-----------------|--|--|--|
| Ref: Bartonella Panel IFA Detects antibodies to Bartonella henselae, Bartonella vinsonii and Bartonella koehlerae in serum via immunofluorescent antibody test. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Testing can be performed on Canine and Feline only. | 60.00 (+ referral fee; see comment) | Referral | M - F | 5-8 business | 2ml Serum | Red top blood collection tube | Refrigerate; send via next day delivery in insulated container with freezer pack(s) |
| Ref: Bartonella PCR This is a polymerase chain reaction assay which amplifies and detects target nucleic acid sequences of the Bartonella genus. This laboratory performs nucleic acid sequencing for all positive samples, verifying detection and providing identification to the species and occasionally strain level. Sample required is EDTA whole blood. This test can be performed on all species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 60.00 (+ referral fee; see comment) | Referral | Tues, Thurs | 5-7 days | 2 mL EDTA whole blood | Purple Top blood Collection Container | Refrigerate, send via overnight delivery; insulated container with freezer pack(s) |
| Ref: Cholinesterase(brain) This assay tests for enzyme activity in a brain sample. Any factor that interferes with enzyme activity will affect the outcome of the results. Specimen should be as fresh as possible Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 48.00 (+ referral fee; see comment) | Referral | M-F | 3-5 days | Brain Tissue/Retina. Split brain into two equal halves sagittally to get a representative hemisphere. Retina-whole eye. | Plastic bag | Keep refrigerated at all times. Ship overnight with ice pack. |
| Ref: Cyclosporine Test to detect cyclosporin levels. This test can be performed on Bovine, Canine, Equine, Feline, Ferrets, Fruit Bats, Ovine, Porcine, Primate samples only. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 66.00 (+ referral fee; see comment) | Referral | W, F | 3-5 business | 1 mL EDTA whole blood | Lavender top blood collection tube | Refrigerate; DO NOT FREEZE. Ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Digoxin Checking Digoxin levels in the blood Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Do not store in rubber topped tubes as digoxin will bind to the rubber. Fasting is recommended, but not required. | 60.00 (+ referral fee; see comment) | Referral | Run when received | 2-4 business | 1 mL serum | red top blood collection tube; GLASS vial | Refrigerate. Ship in insulated container with ice pack. |
| Ref: Feline Calicivirus PCR The Test Target for this Polymerase chain reaction (PCR) assay is the Feline Calicivirus viral RNA, ORF1 Gene This test can be performed on Feline samples only. Submit fresh swabs - do not hold longer than 3 days. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 69.60 (+ referral fee; see comment) | Referral | Mon - Fri | 3 business days | Nasal swab, oropharyngeal swab, nasal biopsy, or lung tissue | Synthetic swab (no cotton tipped swabs) snapped off into a red top tube. Avoid swabs with wooden applicators. Swabs from culturettes are acceptable, but please snap swab off into an empty red top tube. Do not put swab back in transport media/gel. No culture media, viral transport media, or saline solution is necessary. | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s). |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---|------------|-------------|---------------------|--|---|--|
| Ref: Feline Herpes PCR The Test Target is the Viral RNA, GB gene. This test can be performed on Feline samples only. Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 50.00 (+ referral fee; see comment) | Referral | Mon - Fri | 3 business days | Nasal swab, pharyngeal swab, oral swab, or ocular (cornea) swab | synthetic swab (no cotton tipped swabs) snapped off into a red top tube. Avoid swabs with wooden applicators. Swabs from culturettes are acceptable, but please snap swab off into an empty red top tube. Also, avoid putting swab back into media/gel. No culture media, viral transport media, or saline solution is necessary. | Refrigerate. Ship overnight in insulated container with freezer packs. |
| Ref: Feline Infectious Peritonitis mRNA Do not freeze! Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 96.00 (+ referral fee; see comment) | Referral | M-F | 1 day | 2 mL fluid (ascitic, pleural effusions) or EDTA whole blood | | None |
| Ref: Feline Upper Respiratory Disease Panel PCR Includes Chlamydia felis, Feline Calicivirus, Feline Herpesvirus, Bordetella bronchiseptica, Mycoplasma felis. Do not submit swabs in bacterial transport media. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 73.00 (+ referral fee; see comment) | Referral | M-F | 3 days | Deep Pharyngeal swab and Conjunctival swab – plastic-stemmed swabs;no transport media | Red top tube | Refrigerate |
| Ref: FISH Fluorescent In-Situ Hybridization Flourescent probe binds to bacterial ribosomes in tissue. Detected with fluorescent microscope. A \$15 handling fee will be added to each sample, so the total price for a single tissue is \$130. Additional tissues will be billed at \$50 for each additional tissue. For specific information regarding samples, instructions and forms, please see FISH Fluorescent In-Situ Hybridization. Research questions should be directed directly to Simpson Laboratory (607-253-3567). | 115.00 + in-house referral fee. See comment | Referral | Tues or Wed | 5-10 days | 5 unstained fixed tissue slides. If wet fixed tissue or paraffin embedded fixed tissue blocks are submitted, 5 unstained slides per tissue will be prepared for \$25 per tissue. | Slide Mailer; Leakproof, unbreakable container for wet fixed tissue. | Room Temperature; protect wet fixed tissue from freezing |
| Ref: Glomerular Filtration Rate (Iohexol) Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 119.00 (+ referral fee; see comment) | Referral | 1x/wk | 7-10 days | 2 mL serum; samples drawn 2, 3, and 4 hrs after admin of Iohexol | | Refrigerate |
| Ref: Insulin Growth Factor-1 Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Most species can be tested. | 75.60 (+ referral fee; see comment) | Referral | Wednesday | 4 - 9 business days | 0.5 mL Serum or Plasma | Red Top or Purple Top, SST | Refrigerate. Ship in insulated container with cold pack |
| Ref: Keppra/Levetiracetam Checking Keppra levels in the blood Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.Fasting is recommended, but not required. | 84.00 (+ referral fee; see comment) | Referral | Tues, Thurs | 3-4 business days | 1 mL serum | Red top blood collection tube | Refrigerate. Ship in insulated container with ice pack |
| Ref: Leishmania PCR Sample Required is EDTA whole blood. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 60.00 (+ referral fee; see comment) | Referral | M-F | 5-7 days | 2 mL EDTA whole blood | Insulated | Refrigerate or ship on cold pack |
| Ref: Malignancy Profile Profile includes: Parathyroid hormone (PTH), Ionized calcium, Parathyroid hormone related protein (PTHrP) Prefer fasting sample. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 93.60 (+ referral fee; see comment) | Referral | M, Th | 2 - 5 days | 2 mL Serum and 1mL EDTA Plasma | | Refrigerate. Ship in insulated container with freezer packs. |
| Ref: Mycobacterial Culture (Acid Fast TB) Test for presence of tuberculosis. This test does not speciate non-TB Mycobacterium species Send via USPO in NVSL mailer. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.This test is intended to detect Mycobacterium tuberculosis as a reportable disease. | 58.00 (+ referral fee; see comment) | Referral | M-F | 30-120 days | Tissue | Mycobacterium box | Ambient |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|--------------------------------------|------------|-----------|----------|--|---|--|
| Ref: Mycobacterial PCR PCR for Mycobacterium species. Not for M. bovis or M. tuberculosis suspects. Formalin fixed tissue or paraffin blocks should not be decalcified. Tissue should be in wet formalin less than 10 days. Acid fast bacteria must be present on histopathology to perform PCR testing. Fresh tissue should be tested by requesting Mycobacterium Culture rather than PCR to increase testing sensitivity. | 246.00 (+ referral fee; see comment) | Referral | M-F | 12-14 | Busine: Fixed tissue containing lesion | Sterile, leakproof, unbreakable container or paraffin block | Room temperature for fixed tissue and paraffin block |
| Ref: Mycobacterium Culture ID All culture isolates (positive cultures for mycobacterium) will require additional speciation at a fee of \$70.00 to identify the Mycobacteria further. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 48.00 (+ referral fee; see comment) | Referral | Mon - Fri | 60 - 120 | days Fresh tissue | Sterile, leakproof, unbreakable container | Ship for next day delivery in an insulated container with freezer packs. |
| Ref: Pancreatic Lipase Immunoreactivity (PLI), IDEXX Testing for pancreatic inflammatory disease in dogs and cats Send via overnight service; hemolysis and lipemia may interfere with test. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 69.00 (+ referral fee; see comment) | Referral | | 1-2 | days 1 mL CLEAR serum | Insulated | Refrigerate |
| Ref: Parathyroid Hormone Used in diagnosis of: Calcium metabolic disorders, Hypoparathyroidism, Hyperparathyroidism – Primary or Secondary, Malignancy, Vitamin D disorders, Renal disease, Hypercalcemia, Hypocalcemia. Also used to monitor calcitriol therapy. This test can be performed on samples from all species. Send via overnight courier. Avoid Hemolysis and lipemia. Fast animal overnight before taking sample. Ionized Calcium is automatically added to all Parathyroid Hormone requests at no additional charge. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 57.60 (+ referral fee; see comment) | Referral | M-F | 2-6 | Business 1 mL serum only | Red top blood collection tubetube | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Quantitative Calculi/Urinary This test provides an analysis of the crystalline composition of the urolith(urinary stone) Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 38.00 (+ referral fee; see comment) | Referral | M-F | 4-6 | business Stone | Baylor Stone Collection Cup | None |
| Ref: Rabies Titer This test is not used for export testing. This test provides an endpoint titer. Please see comment above regarding export testing. This test can be performed on all species. Send via overnight courier. Can use heat inactivated serum, must be noted on the form. Cannot use plasma. Sample needs at least 500ul for one run. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. The AHDC does not refer Rabies titers for export testing due to requirements of the testing lab. Please contact Kansas State Veterinary Diagnostic Lab directly at 785-532-4483. | 80.04 (+ referral fee; see comment) | Referral | M, Th | 23-30 | Busine: 2 mL serum | Red top blood collection tube | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s). |
| Ref: Rocky Mountain Spotted Fever PCR The sample required is EDTA whole blood. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 60.00 (+ referral fee; see comment) | Referral | M-F | up to 4 | wks 2 mL EDTA whole blood | Insulated | Refrigerate |
| Ref: SDMA Symmetric dimethylarginine (SDMA) is a renal biomarker. This assay which detects SDMA is used to investigate, monitor, and manage kidney disease earlier in dogs and cats. This test is available for Canine and Feline only. Avoid hemolysis and lipemia. Send via overnight courier. Can be stored for 4 days at room temperature, 14 days at 2-8 degrees. Freeze for longer storage. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 12.00 | Referral | M - F | 2 | Business D: 1ml Serum (Preferred) or 1ml EDTA Plasma, or 1 ml Heparinized Plasma | Red Top Collection Container, or Purple Top Collection Container, or Green Top Collection Container | Refrigerate, send via overnight delivery; insulated container with freezer pack(s) |
| Ref: Stone Analysis - Minnesota Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 0.00 (+ referral fee; see comment) | Referral | M-F | 10-14 | days Stone | Unbreakable container | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|--------------------------------------|------------|-----------------|-----------------|---|---|---|
| Ref: Theophylline Checking Theophylline levels in the blood Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Please do not use Serum Separator Tubes. Fasting is recommended, but not required | 60.00 (+ referral fee; see comment) | Referral | Run as received | 2-4 business | 1 mL serum | red top blood collection tube; glass or plastic vial | Refrigerate. Ship in insulated container with ice pack |
| Ref: Tick Panel - PCR PCR Panel includes: PCR testing for known pathogens in six genera: Babesia, Bartonella, Anaplasma, Ehrlichia, Rickettsia, and hemotropic Mycoplasma. All positive PCR results are verified by performing sequencing analysis which also takes identification to the species and sometimes strain level. This test can be performed on most species. Call lab with exotic animals. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 150.00 (+ referral fee; see comment) | Referral | Batched weekly | 9-12 business | 2 mL EDTA whole blood | EDTA blood Tube | Refrigerate, ship for next day delivery in insulated container with freezer pack(s) |
| Ref: TLI, PLI, Cobalamin and Folate Trypsin-Like Immunoreactivity (TLI) assay for exocrine pancreatic function testing (i.e to diagnose EPI) in dogs and cats (species specific) Pancreatic Lipase Immunoreactivity (PLI) assay specifically detects lipase of pancreatic origin for diagnosis of pancreatitis in dogs and cats (species specific). Cobalamin (Vitamin B12) and Folate assay for evaluation of small intestinal function (not species specific) | 86.00 | Referral | M-F | 2-3 Days | 2ml of 12 hour fasted Serum | Red Top Collection Container; send separated serum to avoid hemolysis. | Refrigerate, send via overnight delivery; insulated container with freezer pack(s) |
| Ref: Toxoplasma IgG/IgM This ELISA test detects the presence of IgG and IgM antibodies to Toxoplasma gondii in serum, cerebrospinal fluid and aqueous humor. This test can be performed on Canine, Feline, Canidae, and Felidae only. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 30.00 (+ referral fee; see comment) | Referral | Tu, Thurs | 6 Business days | 0.5 ml Serum, CSF or Aqueous humor | Red Top blood tube | Refrigerate, ship for next day delivery in insulated container with freezer pack(s) |
| Ref: Tritrichomonas foetus PCR Please see male and female sample collection instructions for Tritrichomonas sampling and instructions for using Inpouch mediasystem. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 42.00 (+ referral fee; see comment) | Referral | Tu, Th | 3-5 days | bovine - Tritrichomonas transport media (e.g. InPouch) inoculated with preputial (male) or cervical (female) sample cats - 1-2 grams feces | InPouch media system - bovine samples unbreakable, leakproof tube or cup-feline samples | Room temperature, insulate from temperature extremes |
| Ref: Troponin A complex of muscle proteins which, when combined with Ca ²⁺ , influence tropomyosin to initiate contraction. Send pre and post exercise samples on dry ice by overnight courier. Remove serum or plasma from cells within 2 hours! Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab | 84.00 (+ referral fee; see comment) | Referral | M, F | 1-3 days | 1 mL serum or hep plasma | Insulated | Dry ice |
| Ref: Trypsin-Like Immunoreactivity Exocrine pancreatic function test. This test can be performed on canine and feline samples. Laboratory is willing to run the assay on exotic animal samples, but it is not validated for those. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. | 34.80 (+ referral fee; see comment) | Referral | M-F | 3 - 5 business | 1 mL serum | Red Top blood tube, serum separator tube | Refrigerate, ship for next day delivery in insulated container with freezer pack(s) |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|-------------------------------------|-------------------------|-------------------|--------------|--|--|---|
| Ref: Trypsin-like Immunoreactivity & Pancreatic Lipase Panel | 62.50 (+ referral fee; see comment) | Referral | M-F | 3-5 business | 1 mL serum | Red top blood collection tube or serum separator tube | Refrigerate, ship for next day delivery in insulated container with freezer pack(s) |
| <p>Test for diagnosis of pancreatitis in dogs and cats (species specific) and test for exocrine pancreatic function testing (i.e to diagnose EPI) in dogs and cats (species specific). Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. This test is validated for Canine and Feline. Can be run on Exotics, but it is not validated.</p> | | | | | | | |
| Ref: Vitamin D Profile | 76.80 (+ referral fee; see comment) | Referral | Monday and Friday | 2-8 business | 1.5 mL Serum or Heparinized plasma(not EDTA plasma) | Red Top, Purple Top, Green Top or SST blood tubes | Refrigerate, ship for next day delivery in insulated container with freezer pack(s) |
| <p>Assay for 25- Hydroxyvitamin D This test can be performed on most species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p> | | | | | | | |
| Ref: Vitamin D, 25-hyd | 56.40 (+ referral fee; see comment) | Referral | Monday and Friday | 3-9 Business | 0.5 mL Serum or Plasma (Heparinized and EDTA is accepted) | Red Top, Purple Top, Green Top or SST blood tubes | Red Top, Purple Top, Green Top or SST blood collection tubes |
| <p>1. To identify Vitamin D toxicity or deficiency. A. Will pick up metabolites of cholecalciferol (rat poisons, vitamin D supplements). B. Will not pick up Vitamin D toxicity due to calcitriol (Rocaltrol) or calcipotriene (Dovonex - anti-psoriasis cream). 2. To identify Vitamin D deficiency in animals eating homemade diets, or for certain species, especially Llamas/Alpacas, lizards, primates, and other zoo animals ?This test can be performed on appropriate samples from all species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p> | | | | | | | |
| Ref: Anaplasma phagocytophilum (Ehrlichia Equi) IFA | 37.00 (+ referral fee; see comment) | Referral | M-F | 4-6 business | 2 mL serum | Red top blood collection tube | Refrigerate. Ship for next day delivery in insulated container with freezer pack(s) |
| <p>Detects antibodies against Anaplasma phagocytophilum (formerly known as Ehrlichia equi) in serum via indirect fluorescent antibody test; tested at 1:64, 1:128, 1:256, 1:512, 1:1024, and 1:2048 dilutions. This test can be performed on Equine, Feline, Canine, Ovine, Bovine, Caprine, and Rabbit samples only. Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p> | | | | | | | |
| Reproductive Bacteriology Cultures, Panel 4 | 94.00 /sample | Bacteriology | M-F | 5-7 days | 1) Uterine, cervix or vaginal mucus; 2) Semen, urethra or prepuce | 1-2) Swabs in Amies transport media WITHOUT charcoal or fluids/tissues in sealed sterile container | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| <p>Includes Aerobic, Mycoplasma and Ureaplasma cultures. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not included in the testing fee and are not applicable to Mycoplasma spp. and Ureaplasma spp. Tests can be ordered separately. Preliminary aerobic culture results will be provided one day after sample processing begins.</p> | | | | | | | |
| Reserpine (LC/MS) | 165.50 /sample | Toxicology | M-F | 1-3 days | 1) 10 mL heparin whole blood; or 2) transfer 5 mL heparin plasma to a red top tube | 1) Heparinized tube, no gel clot; 2) red top blood collection tube | 1-2) Refrigerate w/ cool pack |
| <p>Liquid Chromatography/Mass Spectrometry (LC/MS) (increased sensitivity)</p> | | | | | | | |
| Respiratory Bacteriology Cultures, Panel 6 | 94.00 /sample | Bacteriology | M-F | 5 days | 1) Tracheal wash; 2) Lung 3) Aseptically collected swab of tracheal wash or lung | 1-2) Swab in Amies transport media (swab) WITHOUT charcoal or fluids/tissues in sealed sterile container | 1-2) Refrigerate. Ship in insulated container with ice pack. |
| <p>Includes Aerobic, Mycoplasma and Ureaplasma cultures. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not included in the testing fee and are not applicable to Mycoplasma spp. and Ureaplasma spp. Tests can be ordered separately. Preliminary aerobic culture results will be provided one day after sample processing begins.</p> | | | | | | | |
| Reticulocyte Count | 10.50 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| <p>Do not allow coolant to come in contact with specimen; ship Overnight</p> | | | | | | | |
| Rivaroxaban | 45.00 | Comparative Coagulation | As required | Same day | 0.5 mL Citrated plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate sample. Ship overnight with ice packs in insulated container. |
| <p>Anti-Xa activity measured in a chromogenic assay configured with a rivaroxaban calibration standard. Samples for determination of peak rivaroxaban anti-Xa activity should be collected 3 hours after oral drug administration. Frozen plasma will be accepted to analyze.</p> | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---------------|-------------------------|-----------|------------|--|--|---|
| Rotavirus FA-Any Species Fluorescent antibody | 25.00 | Virology | M - F | 2 - 4 days | | Sealable plastic bag or container | Ship in insulated container with ice packs. |
| Russell's Viper Venom Time (RVVT) Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page | 26.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Salmonella Culture For the isolation of Salmonella species. *May qualify for New York State contract pricing. Test is also part of the Enteric Panel 1, Enteric Panel 2, and Enteric Panel 3. Antimicrobial susceptibilities are not included in the testing fee. See the Adult Equine Diarrhea Kit for equine submissions. | 22.00 /sample | Bacteriology | M-Sa | 3-5 days | 1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) other tissues; 5) Water; 6) Swab of 1-4 | 1) Feces in sealed sterile container or ParaPak; 2-5) Tissues or fluids in separate, sealed, sterile containers 6) Swab in Amies transport media or other suitable bacterial transport medium | 1-6) Refrigerate EXCEPT ParaPak (Parapak at room temperature) |
| Selenium, tissue/feed (Selenium Hydride Atomic Absorption) | 55.00 | Toxicology | M-F | 6 days | 1) 2g tissue; 2) 100g feed | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |
| Selenium, whole blood Atomic Absorption analysis of whole blood Cannot analyze in plasma or serum - Need whole blood. Contact Lab for specifics for avian and very small animals. May also want to consider "Vitamin E Alcohol in Serum" test as well. | 28.00 /sample | Toxicology | M-F | 3 days | 1) 5 mL heparin whole blood; 2) 5 mL EDTA whole blood | 1) green top blood collection tube; 2) purple top blood collection tube | 1-2) Refrigerate w/ cool pack |
| Skin Scraping For ectoparasites Visual examination, KOH digest and Sugar Centrifugation Flotation; send an adequate sample size in an escape-proof container; remember to scrape until skin begins to bleed | 28.00 | Parasitology | M-F | 1-3 days | hair/fur and epidermal tissue | escape-proof, sealed container | Not needed |
| Small Animal Chemistry Panel Canine or feline: Na, K, Cl, Bicarb, Anion gap, Na:K, BUN, Creat, Ca, P, Mg, TP, Alb, Glob, A/G, Gluc, ALT, AST, AP, GGT, Tot/Dir/Ind Bilirubin, Chol, CK, LDH, Amylase, Fe, TIBC, % Saturation for additional info on ECLINPATH | 52.00 | Clinical Pathology | M-Sa | 1-2 days | 2 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Small Animal Liver Panel Canine or feline: ALT, AST, ALP, GGT, Tot Bili, Alb, Chol, Gluc, Urea | 27.00 | Clinical Pathology | M-Sa | 4 hours | 1 mL separated serum | plain glass or plastic tube | Refrigerate |
| Small Animal Renal Panel Canine or feline: Na, K, Cl, Bicarb, Anion gap, Urea, Creat, Alb, Ca, P | 28.00 | Clinical Pathology | M-Sa | 4 hours | 1 mL separated serum | plain glass or plastic tube | Refrigerate |
| Sodium (blood, fluid) | 9.00 /sample | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| Sodium, urine | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| Sorbitol Dehydrogenase (SDH), blood | 11.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Special Stain #1 THIS TEST CODE FOR INTERNAL USE ONLY. | 50.00 | Clinical Pathology | M - F | 5 days | Slides | | |
| Special Stain #2 | 80.00 | Clinical Pathology | M-F | 5 days | Slides | | |
| Special Stain #3 | 110.00 | Clinical Pathology | M - F | 5 days | SLIDES | | |
| Special Stain #4 | 140.00 | Clinical Pathology | M - F | 5 days | Slides | | |

Last Updated: 4/3/2017
VSS-WEB-082-V01

* Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 31 of 36

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---------------|--------------------|-----------|-----------------|--|---|--|
| Stained Histology Slides Slides prepared from (FFPE) blocks H&E or Special Stained slides prepared from Anatomic Pathology cases in addition to the Histopathology. Stained slides are requested to either have Anatomic Pathology review or to have them sent to another facility for review/referral. Additional charges will be applied for shipping and handling. | 10.25 | Anatomic Pathology | M-F | 5 business days | Paraffin embedded tissue block | | Room temperature |
| Strychnine, liquids/fluids | 55.25 /sample | Toxicology | M-F | 3 days | 1) 10 - 20 mL heparin whole blood; 2) 10 - 20 mL urine | 1) Heparinized tube, no gel clot; 2) sterile leak-proof container | 1-2) Refrigerate w/ cool pack |
| Strychnine, solids | 90.00 /sample | Toxicology | M-F | 3 days | 1) 50g liver; 2) 50g stomach contents or vomitus | 1-2) sterile container | 1-2) Refrigerate w/ cool pack |
| Sulfur | 40.00 | Toxicology | M-F | 3 days | 1) 2g tissue; 2) 100g feed | 1) sterile container; 2) any container | 1) Refrigerate; 2) none |
| Synovial Fluid Cytology Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Viscosity, Smear type, Smear description, Smear Interpretation If only slides (and no fluid) are submitted, request a cytology smear exam on the cytology submission form and specify the joint sampled as the source. For joint fluid in EDTA, please also submit 2-3 freshly prepared smears from the fluid. Also specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. It is important to rapidly air-dry any smears. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | 55.00 | Clinical Pathology | M - F | 1 business day | 1) Joint fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment. |
| T3 (Triiodothyronine) baseline For more information, see Appendix C | 17.00 | Endocrinology | M-F | 1-3 days | 1.5 mL serum or heparinized plasma | plastic or glass tube | Refrigerate |
| T3 Post Suppression Test: T3 Contact Lab for protocol; For more information, see Appendix C | 11.50 | Endocrinology | M-F | 1-3 days | 2 mL serum or heparinized plasma | plastic or glass tube | Refrigerate |
| T3 Post Suppression Test: T4 Contact Lab for protocol; For more information, see Appendix C | 10.50 | Endocrinology | M-F | 1-3 days | 2 mL serum or heparinized plasma | plastic or glass tube | Refrigerate |
| T3 Suppression Test: T3 T3; feline Contact Lab for protocol; For more information, see Appendix C | 15.50 | Endocrinology | M-F | 1-3 days | 2 mL serum or heparinized plasma | plastic or glass tube | Refrigerate |
| T3 Suppression Test: T4 Contact Lab for protocol; For more information, see Appendix C | 13.50 | Endocrinology | M-F | 1-3 days | 2 mL serum or heparinized plasma each sample | plastic or glass tube | Refrigerate |
| T4 (Thyroxine) baseline Thyroxine baseline For more information, see Appendix C | 15.00 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate |
| T4 & Free-T4 by dialysis & TSH combination Includes T4, FT4d, TSH For more information, see Appendix C | 58.00 | Endocrinology | M, Th | 2-5 days | 2 - 2.5 mL serum | red top blood collection tube or plastic tube | Refrigerate |
| T4 & Free-T4 by dialysis combination Includes T4 and FT4d For more information, see Appendix C | 47.00 | Endocrinology | M, Th | 1-5 days | 1 - 2 mL serum | plastic or glass tube | Refrigerate |
| T4 & TSH combination Includes T4 and TSH; canine, feline For more information, see Appendix C | 28.00 | Endocrinology | M-F | 1 day | 1 - 2 mL serum | plastic or glass tube | Refrigerate |
| T4T3 This assay used for small animal testing | 27.00 | Endocrinology | M-F | 1-3 days | 400 uL serum | red top blood tube | Refrigerate |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|----------------------|-------------------------|-----------|---------------|--|--|--|
| Testosterone baseline For more information, see Appendix C | 19.00 | Endocrinology | Monday | 1-8 days | 1 mL serum | plastic or glass tube | Refrigerate |
| Testosterone Post HCG/GnRH | 13.00 | Endocrinology | Monday | 1-7 days | 250 uL serum or plasma | plastic or glass | |
| Testosterone: HCG/GnRH Response | 19.00 | Endocrinology | Monday | 1-7 days | 250 uL serum or plasma | plastic or glass | |
| Thrombin Clotting Time (TCT) Ship Overnight: for additional info on Submission Guidelines, or see Comp Coag web page | 14.50 | Comparative Coagulation | M-F | same day | 1 mL separated citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| Thrombin-antithrombin Complex Research and batch testing only. Testing not done on rabbits. Per test fee requires submission of 15 or more samples for batch analysis. Ship Overnight. For additional info on Submission Guidelines, or see Comp Coag web page: | 52.50 | Comparative Coagulation | 1x/month | up to 1 month | citrate plasma | plastic tube in styrofoam box | dry ice |
| Thyroid Panel, Feline: FT4d, T4, T3, TSH For more information, see Appendix C | 62.00 | Endocrinology | M, W, Th | 2-5 days | 2 - 2.5 mL serum or heparinized plasma from a plain red top collection tube, with no additives and no serum separator tubes; should arrive separated and chilled | red top blood collection tube, plastic or polypropylene tube adequate for transport and frozen storage | Refrigerate |
| Thyroid Panel: FT4 by dialysis, T4, T3 For more information, see Appendix C | 57.00 | Endocrinology | M, Th | 1-5 days | 2 mL serum or heparinized plasma | plastic or glass tube | Refrigerate |
| Total Protein Panel Total Protein (TP), Albumin, Globulin for additional info on Clin Path panels | 17.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Total Protein Plasma (by refractometer) Do not allow coolant to come in contact with specimen; ship Overnight | 5.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (>1/2 full) | lavender top tube | Refrigerate |
| Total Protein, blood Total protein in blood | 13.00 | Clinical Pathology | M-Sa | 1-2 days | 1 mL separated serum or heparinized plasma | plain glass or plastic tube | Refrigerate |
| Total Protein-Creatinine Panel Total Protein, Creatinine, P:C Ratio (Calc) for additional info on Clin Path panels | 38.00 | Clinical Pathology | M-Sa | 1-2 days | 2 mL urine | leak-proof container | Refrigerate |
| Toxic Heavy Metal Screen (Arsenic, Lead, Cadmium, Mercury, Thallium) X-ray Fluorescence Spectrophotometry | 77.25 /sample | Toxicology | M-F | 7 days | 1) liver OR 2) kidney OR 3) powders, solids OR 4) paint chips | 1-5) sterile, leak-proof container | 1-2) Freeze; 3-5) Refrigerate w/ cool pack |
| Toxoplasma gondii IgG ELISA Detects anti-Toxoplasma IgG antibody The Toxoplasma gondii IgG ELISA tests serum for the presence of anti-Toxoplasma IgG antibody, indicating a history of exposure to the parasite. This test is valid to test serum from a broad range of animal species, except for birds, marsupials and cold-blooded animals. Results are reported as "positive", "negative" or "equivocal". Testing is repeated 2-3 weeks after the initial sample was tested to establish if an infection is acute or chronic, or to delineate the true status of an animal with a previously "equivocal" result. When a second sample is submitted, please provide the accession number from the first sample submitted, as both samples are needed for accurate paired testing. Fluids other than serum will not be evaluated. *May qualify for New York State contract pricing. | 20.00 /sample price. | Parasitology | Wed | 1-7 days | 100 uL serum | red-top blood collection tube | Refrigerate |
| Toxoplasmosis FA Fluorescent antibody | 25.00 /animal | Virology | M-F | 2-4 days | Fetal tissue: placenta and brain preferred; also lung, liver, kidney, skeletal muscle, spleen, lymph node; Non-fetal tissue including suspect affected tissue such as brain, liver, kidney, lung, skeletal muscle. | Leakproof, unbreakable plastic container or bag | Ship for next day delivery in insulated container with freezer pack(s) |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---------------------------------|--------------------|-----------|-----------------|---|--|--|
| Tracheal Wash Cytology | 46.00 | Clinical Pathology | M - F | 1 business day | 1) Tracheal wash fluid in EDTA; 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated. | 1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides | Refrigerate fluids only, DO NOT refrigerate smears |
| Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation With fluid samples of tracheal wash, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. It is important to rapidly air-dry any smears. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing | | | | | | | |
| Trichinella Tissue Examination Acid Pepsin digest | 27.50 | Parasitology | M-F | 7 days | muscle tissue | wide mouth leak-proof | Refrigerate |
| Trichomonas Fecal Examination | 25.00 | Parasitology | M-F | 12 days | fecal swab in trichomonas In-Pouch (see instructions with In-Pouch) | trichomonas In-Pouch | Do NOT Refrigerate |
| Trichomonas pouches are available from AHDC Shipping | | | | | | | |
| Triglycerides | 11.00 /sample (blood, fluid) | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate |
| TSH (Thyroid Stimulating Hormone) baseline Canine, feline For more information, see Appendix C | 19.00 | Endocrinology | M-F | 1-3 days | 1 mL serum | plastic or glass tube | Refrigerate |
| Unstained Histology Slides Prepared | 5.25 | Anatomic Pathology | M-F | 5 business days | Paraffin embedded tissue block | | Room temperature |
| Slides or PCR tubes prepared from (FFPE) blocks. Unstained slides or PCR tubes containing 2-3 (10uM) unstained tissue sections prepared from anatomic pathology cases that have been or are in the process of being reported on. These unstained slides or PCR tubes are prepared for the purpose of sending to another testing facility for slide review or referral analysis. Additional charges may be applied for shipping and handling. Price is per slide or PCR tube prepared. | | | | | | | |
| Urea Nitrogen | 11.00 | Clinical Pathology | M-Sa | 1-2 days | 1) 1 mL separated serum or heparinized plasma; 2) fluid | plain glass or plastic tube | Refrigerate. Ship in insulated container with ice pack. |
| Test for blood urea nitrogen in serum, plasma or other fluid. | | | | | | | |
| Urea Nitrogen, urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate. Ship in insulated container with ice pack. |
| Ureaplasma Culture | 27.00 /sample | Bacteriology | M-F | 5-10 days | 1) Uterine, cervical or vaginal mucus; 2) Semen; 3) Tracheal washing or lung tissue; 4) Urine; 5) Aseptically collected swab of 1-4 | 1-4) Fluid or tissue in sealed sterile container; 5) Swab in Amies transport media WITHOUT charcoal | 1-4) Refrigerate |
| For the culture of Ureaplasma species. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Test is also part of the Reproductive Panel 4 and Respiratory Panel 6. | | | | | | | |
| Uric Acid, urine | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1 mL urine | plain glass or plastic tube | Refrigerate |
| Urinalysis, Routine | 25.00 | Clinical Pathology | M-Sa | 1-2 days | 10 mL fresh urine | leakproof container | Refrigerate |
| Color, turbidity, dipstick (pH, TP, glucose, ketones, bilirubin, blood), specific gravity, sediment exam | | | | | | | |
| Urine Culture | 20.00 /sample | Bacteriology | M-F | 2-7 days | 1) Urine; 2) Bladder tissue/stones; 3) Aseptically collected Swab of urine or bladder | 1) Urine Transport Tube (preferred) or other sterile container; 2) Tissue or stones in separate sterile sealed containers; 3) Swab in Amies transport media or other suitable transport medium for aerobic culture | (1-3) Ship overnight in insulated container with ice packs to arrive no later than 48 hours post collection. |
| For the isolation of aerobic bacteria from the urinary tract. *May qualify for New York State contract pricing. It is recommended to collect and submit urine collected by Cystocentesis. Please indicate method of collection. Antimicrobial susceptibilities are not included in the testing fee. Refer to Urine Antimicrobial susceptibilities. Urine Transport Tubes are available for purchase from our Supply Center. Performing cytology in conjunction with culture is also recommended to aid in interpretation of results. | | | | | | | |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|--|---------------|-------------------------|------------------|---------------|---|--|---|
| Virus FA See also specific Viral FA tests Refer to individual FA tests | 25.00 /animal | Virology | M-F | 2-4 days | Tissues or fluids. Contact lab if there are questions about sample suitability. | Sealable plastic bag or container for tissues; Leakproof container for fluids and swabs in Viral Transport Medium | Ship in insulated container with ice pack |
| Virus Isolation See also specific Virus Isolation tests or clinical presentation Best sample may depend upon suspected viral infection and pathophysiology related to viremia, infected tissue and viral shedding. Recent investigations indicate that urine is a good antemortem sample for detection of some viruses. There is generally a lack of information regarding viral detection in urine samples for many viral diseases. Please include urine if it is available when you are submitting other samples for virus isolation for our investigational use. No extra charges will be incurred. Refer also to individual Virus Isolation tests. Call lab ahead before sending reptile samples. Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline. | 70.00 /animal | Virology | M-F | 21-30 days, s | 1) Tissue [target organs and blood-rich organs such as lymph nodes, spleen, lungs, kidney, heart, liver, brain] OR 2) Cerebrospinal fluid Or 3) EDTA whole blood OR 4) antemortem nasal swab AND 5) urine (see comment) | 1) Tissue in leakproof unbreakable containers 2) sterile, leakproof container; 3) EDTA blood tube; 4) in sterile leakproof container with 2-3 drops sterile saline or in Virus isolation media (not bacterial transport media) | Ship in insulated container with freezer packs for next day delivery pack |
| Vitamin A in serum (High Performance Liquid Chromatography) Analysis for retinol (vitamin A) using HPLC. Separate serum and remove from clot immediately. Protect from light and keep cool. | 30.00 each | Toxicology | Tu, Th | 5 days | Serum, separated from clot. | Avoid preservatives such as Citrate or EDTA. No Gel Clot Vacutainer Tubes. | Refrigerate |
| Vitamin B12 & Folate combination Avoid hemolysis and excessive exposure to direct light. For more information, see Appendix C | 40.00 | Endocrinology | Tu | 1-8 days | 1.5 - 2 mL serum; avoid EDTA plasma | plastic or glass tube | Refrigerate. Ship in insulated container with ice pack. |
| Vitamin B12, Cobalamine Vitamin B12 Avoid hemolysis and excessive exposure to direct light. For more information, see Appendix C | 23.00 | Endocrinology | Tu | 1-8 days | 1 mL serum; avoid EDTA | plastic or glass tube | Refrigerate. Ship in insulated container with ice pack. |
| Vitamin E in serum (High Performance Liquid Chromatography) "Selenium" test recommended also Analysis for alpha-tocopherol (vitamin E) using HPLC. Separate serum and remove from clot immediately. Failure to remove serum from clot may lead to inaccurately low vitamin E levels. Protect from light and keep cool. We strongly recommend that Vit E (serum) and Selenium (Heparin or EDTA whole blood) be tested together. | 30.00 each | Toxicology | Tu, Th | 5 days | Serum, separated from clot | Avoid preservatives such as Citrate or EDTA. No Gel Clot Vacutainer Tubes (SST tubes). | Refrigerate |
| Vitamin K Deficiency Coag Panel Incl. aPTT, PT, TCT, Factor VII:C to detect vitamin K-dependent coagulopathy including anticoagulant rodenticide toxicity. Panel for dogs and cats only, see Vitamin K Deficiency Coagulation Panel FAQs and Vitamin K Dependent Coagulopathy. | 41.50 | Comparative Coagulation | M - F | 24 hours | 2 mL citrated plasma | sodium citrate anticoagulant (blue top tube) | Freeze. Ship in insulated container with ice packs |
| von Willebrand factor (vWF:Ag) von Willebrand factor concentration, ELISA method Ship Overnight by FedEx or UPS; for additional info on Submission Guidelines, or see Comp Coag web page | 32.00 each | Comparative Coagulation | Tues, Thurs, Fri | 1-3 days; San | 1 mL separated citrate plasma or EDTA plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |
| von Willebrand factor Collagen Binding Assay Functional ELISA Ship Overnight; requires vWF:Ag on same sample; for additional info on Submission Guidelines, or see Comp Coag web page | 43.00 | Comparative Coagulation | bi-weekly | 1-2 wks | citrate plasma | plastic or siliconized glass tube in styrofoam box | Refrigerate/ship on cold pack |

Animal Health Diagnostic Center Test and Fee Schedule

| Test Name | Test Fee | Discipline | Test Days | Lag** | Samples | Container | Coolant |
|---|---------------|--------------------|-----------|-----------|---|---|---|
| West Nile Virus (WNV) PCR PCR Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. Whole blood is an acceptable sample only for Avian testing, not for Mammals This service is performed pursuant to an agreement with Roche Molecular Systems. | 36.75 | Virology | M-F | 3-5 days | AVIAN: 1) Brain, heart, kidney, bone marrow; 2) EDTA Whole Blood; 3) Oropharyngeal or Cloacal swab; MAMMAL:4) CSF; 5) Brain or spinal cord. WHOLE BLOOD IS NOT AN ACCEPTABLE SPECIMEN FOR MAMMALS | 1,5) Sealable plastic bags or containers; 3) Swabs in Virus transport medium; 2,4) Leakproof container | Freezer Packs |
| West Nile Virus (WNV) SN serum neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Acute and convalescent serum samples are recommended for optimal diagnostic test interpretation. Currently at the AHDC, this test is used primarily for mammalian species. | 14.50 | Virology | Thursday | 7-14 days | 1 ml of serum (preferred) or plasma | red top blood collection tube | It is recommended to ship samples with an ice pack. |
| White Blood Cell (WBC) Count, Total Do not allow coolant to come in contact with specimen; ship Overnight | 7.00 | Clinical Pathology | M-Sa | 1-2 days | EDTA blood (> 1/2 full) | lavender top tube | Refrigerate |
| White Blood Cell (WBC) Panel WBC, Manual Differential Cell Count, Platelet Smear estimate, RBC, WBC Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panels | 16.50 | Clinical Pathology | M-Sa | 1-2 days | 1) EDTA blood > 1/2 full; 2) 2 dry (unfixed, unstained) blood smears | 1) lavender top tube; 2) frosted-end slides | Refrigerate. Ship in insulated container with ice packs. |
| Yersinia Culture For the isolation of Yersinia enterocolitica. Antimicrobial susceptibilities are not included in the testing fee. | 16.00 /sample | Bacteriology | M-Sa | 2-7 days | 1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) swab of 1-3 | 1) Feces in sealed sterile container or ParaPak; 2-3) Tissue in Separate sealed sterile containers; 4) Swab in Amies transport media or other suitable aerobic transport medium | 1-3) Refrigerate EXCEPT ParaPak (keep at room temperature) |
| Zinc, serum/plasma For avian and exotics, contact Lab | 30.00 /sample | Toxicology | M-F | 2 days | 1) 2 mL serum; or 2) 2 mL heparinized plasma in a green top tube | 1) 2 mL serum; or 2) 2 mL heparinized plasma; Royal blue-topped tube (trace mineral tube) preferred. No gel clot. | 1-2) Refrigerate. Ship in insulated container with ice packs. |