

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Acid Fast Stain (for bacteria)	8.50 /sample	Bacteriology	M-F	1-2 days	1) Affected tissue, lesion, or fluid; 2) Impression smear	1) Amies transport media (swab) or sealed sterile container; 2) slide mailer	1) Refrigerate. Ship in insulated container with ice pack. 2) Not required
For the direct microscopic detection of Mycobacterium species. *May qualify for New York State contract pricing							
Acidic-Extracted Drugs	70.00 STAT+25	Toxicology	M-F	2 business days	1) 4 ml heparinized (preferred) or EDTA plasma or serum; OR 2) 10 ml heparinized (preferred) or EDTA whole blood; OR 3) 50 ml urine	Heparinized tube(s), no gel clot; or red top tube with transferred heparinized plasma, labeled accordingly. Less ideal: red top tube with transferred serum or EDTA plasma OR red top/lavender top tube(s) with whole blood OR urine in leak-proof container	1-3) Refrigerate. Ship for next day delivery in insulated container with freezer pack(s)
Qualitative drug screening and confirmation. Analytes include, but are not limited to: Acetaminophen, Barbiturates, Bumetanide, Chlorothiazide, Diclofenac, Diflunisal, Firocoxib, Flufenamic Acid, Flunixin, Flurbiprofen, Furosemide, Griseofulvin, Ibuprofen, Indomethacin, Ketoprofen, Meclofenamic Acid, Meprobamate, Methocarbamol, Naproxen, Oxyphenbutazone, Phenylbutazone, Nabumetone, Salicylic Acid, Sulfa Drugs (Sul diazine, Sulfadimethoxine, Sulfadimidine, Sulfafurazole, Sulfamerazine, Sulfamethoxazole, Sulfanilamide, Sulfapyridine, Sulfathiazole), Sulindac, Thiosalicylic Acid, Tolmetin, Zomepirac.							
Aerobic Bacterial Culture	40.00 /sample	Bacteriology	M-F	2-7 days	Affected tissue, lesion, fluid aspirate or swab in bacterial transport media	Amies transport media (swab) or sealed sterile container	Refrigerate. Ship in insulated container with ice pack.
For the isolation of aerobic bacteria. Tissues are not routinely pooled for aerobic cultures. When multiple tissues are submitted and aerobic culture is requested, we will culture each separately, and a fee for each tissue will accrue. To detect Salmonella from GI samples, a Salmonella culture should be requested. Salmonella is readily identified in aerobic cultures from non-GI sources. To detect Listeria in brain or CSF specimens, a Listeria culture should be requested. Listeria is readily identified in aerobic cultures from systemic samples of septicemic animals or fetal tissues. Streptococcus equi is identified in routine aerobic cultures. If no other aerobic bacteria are of diagnostic concern, a Strep. equi culture can be used to rule in/out Strep. equi only. For Campylobacter spp., Mycoplasma spp., Ureaplasma spp., Yersinia spp. and Brucella spp., individual cultures specific to the bacteria should be requested, as they are not detected reliably in routine aerobic cultures. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not included in the testing fee. Susceptibilities will be performed only if specifically requested. See Antimicrobial Susceptibility (MIC) and Antimicrobial Susceptibility, Ophthalmology. Preliminary results will be provided one day after sample processing begins.							
Alanine Aminotransferase (ALT)	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
See Eclinpath for more information							
Albumin	8.50	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship in insulated container with ice pack.
See Eclinpath for more information							
Alkaline Phosphatase (ALP)	8.50 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid;	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
See Eclinpath for more information							
Alkaloid Screen	331.00 /sample	Toxicology	M-F, as needed	5-7 days	1) liver; 2) heparin whole blood; 3) urine; 4) other materials	1-3) unbreakable sterile container; 4) Heparinized tube, no gel clot	1-4) Refrigerate. Ship in insulated container with ice pack.
Qualitative screening for alkaloids such as strychnine, brucine, caffeine, theobromine, theophylline, nicotine, atropine, cocaine, morphine, codeine, ephedrine, reserpine, quinine, and solanidine, on request. Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis.							
Ammonia	54.00	Clinical Pathology	M-F	1 day	Separated plasma from a purple top (EDTA).	Red top (with separated plasma) or purple top (whole blood)	Refrigerate: 4 C or frozen (on dry ice) – see Appendix A for special handling instructions
Used to identify hyperammonemia in cases with neurologic symptoms due to hepatic failure (hepatic encephalopathy), enteric disease (colonic disorders in horses) and inherited disorders of ammonia metabolism. Call the lab at 607-253-3255 prior to drawing sample. See Eclinpath for more information							
Amylase	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship in insulated container with ice pack.
See Eclinpath for more information							

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Amylase, urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer pack.
Anabolic Steroids in Urine	441.00 /sample	Toxicology	M-F as needed	7 business days	urine or plasma	sterile container	Refrigerate. Ship in insulated container with ice pack.
Qualitative screening for boldenone, dianabol, and stanozolol. EFFECTIVE 6/7/17, ANABOLIC STEROID TESTING IS CURRENTLY UNAVAILABLE. SUBMITTED SAMPLES WILL BE FORWARDED TO A REFERRAL LAB FOR ANALYSIS.							
Anaerobic Bacterial Culture	45.00 /sample	Bacteriology	M-F	2-7 days	1) Fluid aspirate inoculated into anaerobic transport media vials for liquid specimens(Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM); 2) Fresh tissue (>2-3 cm in diameter, minimum size); 3) Ligated loop of intestine; 4) Swab or small tissue collected and immediately placed in anaerobic transport media tube for swabs(Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM)	1,4) Anaerobic transport media (Cary-Blair, e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM): tube or vial form - see sample info above; 2-3) tissues (>2-3 cm in diameter, minimum size)(not swabs) in sealed sterile container	1) DO NOT Refrigerate anaerobic transport media tubes or vials; 2-3) Tissues (>2-3 cm in diameter, minimum size) are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection.
For the isolation of anaerobic bacteria. Ship overnight. Antimicrobial susceptibilities are not applicable. Please consider using the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine diarrhea submissions. Tissues (>2-3 cm in diameter, minimum size or ligated loops of bowel) are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection. DO NOT Refrigerate anaerobic transport media tubes or vials Despite labeling to the contrary, culture systems containing Amies transport media or Stuart transport media are inappropriate as transport systems for anaerobic culturing. Swabs transported without any transport media are unsuitable for both aerobic and anaerobic culture.							
Anesthetic Screen	331.00 per /sample	Toxicology	M-F as needed	3 business days	1) urine; 2) heparin whole blood; 3) liver	1) sterile, leak-proof container; 2) Heparinized tube; 3) sterile, leak-proof container	1-3) Refrigerate. Ship in insulated container with ice pack.
Qualitative screening for acepromazine, barbiturates, ketamine, morphine, xylazine, tiletamine/zolazepam. Special request only; contact Lab.							
Anticoag/Rodenticide Screen, liquid	165.50 /sample	Toxicology	M-F	3 business days	1) heparin whole blood; 2) urine	1) Heparinized tube; 2) sterile container	1-2) Refrigerate. Ship in insulated container with ice pack.
Screening for brodifacoum, bromadiolone, chlorophacinone, dicoumarol, difenacoum, diphacinone, flocoumafen, warfarin For more information, see Strychnine, liquids/fluids.							
Anticoagulant/Rodenticide Screen, solids	165.50 /sample	Toxicology	M-F	5 business days	1) liver; 2) stomach contents; 3) feed	1-3) sterile container	1-2) Freeze. Ship in insulated containers with ice packs.
Qualitative testing for Brodifacoum, Bromadiolone, Chlorophacinone, Dicoumarol, Difenacoum, Diphacinone, Flocoumafen, Warfarin For more information, see Strychnine, solids.							
Antimicrobial Susceptibility (MIC)	28.00 /isolate	Bacteriology	M-F	2-5 days	1) Affected tissue, lesion or fluid; 2) Feces or intestinal content; 3) Bacterial isolate	1) Amies transport media (swab) or other appropriate bacterial transport media OR 2) Exudate or aspirate fluid in sealed sterile container OR 3) Bacterial isolate on Agar slant or plate (sealed) or swab in appropriate bacterial transport media	1-3) Refrigerate. Ship in insulated container with ice pack.
Includes systemic antimicrobials. Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. **May qualify for New York State contract pricing .							

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Antimicrobial Susceptibility, Ophthalmology	30.50 per isolate	Bacteriology	M-F	2-5 days	1) Bacterial isolate	1) In house culture isolate 2) Agar slant or plate (sealed)	1-2) Refrigerate. Ship in insulated container with ice pack.
Includes topical ophthalmic antimicrobials. *May qualify for New York State contract pricing . Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. Special certified training is required for the shipment of some pure cultures. Contact AHDC laboratory with questions							
Antimicrobial Susceptibility, Urine	24.50 /isolate	Bacteriology	M-F	2-5 days	1) Urine; 2) Bladder tissue/stones; 3) Bacterial isolate** (see Comment)	1-2) Sterile container or Amies transport media (swab); 3) Agar slant or plate (sealed)	1-3) Refrigerate. Ship in insulated container with ice packs.
Includes antimicrobials that concentrate in the bladder. *May qualify for New York State contract pricing . Bacterial culture must be performed to recover an isolate before testing can be done. Cost of culture is separate. For susceptibilities on bacterial isolates, pure, identified cultures must be submitted. **Shippers of bacterial isolates must be fully trained in Hazardous Materials Shipping and adhere to all applicable regulations.							
Antiplasmin	38.50	Comparative Coagulation	M-F	same day	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate. Ship in insulated container with ice pack.
Measurement of antiplasmin levels in plasma Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page							
Antithrombin (AT3)	35.75	Comparative Coagulation	M-F	same day	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate. Ship in insulated container with ice pack.
Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page							
aPTT (Activated Partial Thromboplastin Time)	14.50	Comparative Coagulation	M-F	same day	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate. Ship in insulated container with ice pack.
Ship Overnight. For additional info see Submission Guidelines, or see Comp Coag web page							
aPTT Inhibitor Screen	14.50	Comparative Coagulation	M-F	same day	citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate. Ship in insulated container with ice pack.
Ship Overnight; requires aPTT on same sample; for additional info see Submission Guidelines, or Comp Coag web page							
Arsenic, blood/water/urine	40.00 /sample	Toxicology	M-F	3 business d	1) whole blood; 2) urine; 3) water	1) Heparinized or EDTA tube; 2) urine in sterile container	1-2) Refrigerate. Ship in insulated container with ice pack.
Quantitative arsenic concentration in fluid samples. Cannot analyze in plasma or serum - Need whole blood. Do not use heparin tubes (gel tubes). Whole blood should not be clotted. Contact Lab for specifics for avian and very small animals							
Arsenic, tissue/feed	35.00 /sample	Toxicology	M-F	5 business d	1) tissue; 2) feed	1) sterile container; 2) any container	1) Refrigerate; 2) none
Quantitative arsenic concentration in tissue or feed.							
Aspartate Aminotransferase (AST)	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
Aspartate Aminotransferase (AST), urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate. Ship in insulated container with ice pack.
Bacterial ID	40.00 /isolate	Bacteriology	M-F	2-7 days	Bacterial isolate (pure culture)	Agar slant or plate (sealed) or Amies transport media (swab)	Refrigerate. Ship in insulated container with ice pack.
For the identification of pure bacterial isolates. Charges are assessed PER ISOLATE submitted. Please ensure purity before shipment. Submissions accepted from Laboratories only. Follow all biohazard shipping regulations. Please provide animal history and suspected identification of isolate if possible.							

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Baermann Fecal Technique	25.00	Parasitology	M-F	2-3 days	15g fresh fecal sample (~1/2 cup)	Plastic leakproof container	Refrigerate. Ship in insulated container with ice pack.
<p>A modification of the Berlese Apparatus used to retrieve nematode such as lungworm, larvae from feces. The Baermann Technique is not recommended as a primary diagnostic technique for evaluation of parasites in feces. It is not useful for those nematode larvae that do not leave the feces or other biological material, or for detection of parasite eggs or cysts that may be in the fecal sample. Some lungworm larvae are retrieved using this technique, but some larvae such as those of <i>Filaroides hirithi</i>, <i>Filaroides osleri</i>, <i>Strongyloides</i> sp. and <i>Dictyocaulus</i> sp. are better recovered using flotation techniques. Lungworms, such as <i>Eucolus</i> (<i>Capillaria</i>) <i>aerophilus</i>, that do not produce larvae are detected only by using flotation methods as well. Any fecal sample submitted for this procedure must be freshly voided so that the sample is not contaminated by free-living nematodes. It is strongly recommended to use a fecal flotation technique when lungworms are suspected.</p>							
Basic-Extracted Drugs	70.00 /sample; plus \$25 for STAT	Toxicology	M-F	2 days	1) 4 ml heparinized (preferred) or EDTA plasma or serum; OR 2) 10 ml heparinized (preferred) or EDTA whole blood; OR 3) 50 ml urine	Heparinized tube(s), no gel clot; or red top tube with transferred heparinized plasma, labeled accordingly. Less ideal: red top tube with transferred serum or EDTA plasma OR red top/lavender top tube(s) with whole blood OR urine in leak-proof container	1-3) Refrigerate. Ship for next day delivery in insulated container with freezer pack(s)
<p>(including but not limited to): Acepromazine, Amphetamine, Apomorphine, Atropine, Benzocaine, Brompheniramine, Brucine, Bupivacaine, Caffeine, Cimetidine (Tagamet), Chlorpromazine, Cocaine, Codeine, Cyproheptadine, Dextromethorphan, Diazepam, Dihydrocodeine, Dipyrone, Doxapram, Embutramide, Ephedrine, Griseofulvin, Hordenine, Hydrocodone, Ketamine, Lidocaine, Mepivacaine, Meprobamate, Methocarbamol, Methylphenidate, Methamphetamine, Metoclopramide, Pentazocine, Phenacetin, Phentoiazine, Procaine, Promazine, Phenylpropanolamine, Propoxyphene, Pyrilamine, Rantidine (Zantac), Strychnine, Tetracaine, Theobromine, Theophylline, Trimethoprim, Tropicamide, Xylazine For more information, see Appendix B. \$25.00 will be added for STAT processing.</p>							
Beta-Hydroxybutyrate (BHBA)	13.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum, EDTA plasma, or heparinized plasma	Red top (non-anticoagulant), green top (heparin) tube, or purple top (EDTA tube)	Refrigerate. Ship overnight in insulated container with freezer pack.
<p>See Eclinpath for more information</p>							
Bethesda Unit Assay	52.00	Comparative Coagulation	M	1 wk	citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate. Ship in insulated container with ice pack.
<p>Bethesda units/mL: measure of factor inhibitory activity Ship Overnight. For additional info see Submission Guidelines or Comp Coag web page</p>							
Bicarbonate	9.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
<p>See Eclinpath for more information</p>							
Bile Acids, Fasting Panel	24.00 /sample	Clinical Pathology	M-Sa	1-2 days	1 - 2 mL separated serum, each sample	Red top (non-anticoagulant) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
<p>Fasting Panel Clearly mark "Fasting" on sample. Lipemia (falsely increases) and hemolysis (falsely decreases) interfere with bile acid measurement through optical interference. Efforts should be made to minimize these (by not feeding too large a meal and by separating serum from cells as soon as possible). See Eclinpath for more information</p>							
Bile Acids, Post-Prandial Panel	24.00 /sample	Clinical Pathology	M-Sa	1-2 days	1 - 2 mL separated serum, each sample	plain glass or plastic tube	Refrigerate. Ship overnight in insulated container with freezer packs.
<p>Post-Prandial Panel</p>							
Bile Acids, Random Panel	24.00 /sample	Clinical Pathology	M-Sa	1-2 days	1 - 2 mL separated serum, each sample	Red top (non-anticoagulant) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
<p>Lipemia (falsely increases) and hemolysis (falsely decreases) interfere with bile acid measurement through optical interference. Efforts should be made to minimize these (by not feeding too large a meal and by separating serum from cells as soon as possible).</p>							
Bilirubin Panel (total, direct, indirect)	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship in insulated container with ice pack.
<p>Includes Total, direct, and indirect bilirubin. See Eclinpath for more information</p>							

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Billirubin, Total See Eclinpath for more information	8.50 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid;	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Blood Culture, Aerobic For the isolation of aerobic bacteria in blood. Blood culture medium should be inoculated immediately when blood is drawn. See https://ahdc.vet.cornell.edu/docs/Blood_Culture_Technique.pdf for instructions. If the blood culture media used is not the VersaTrek EZ® Draw® REDOX 1® product, turn around time may be delayed due to inability to take full advantage of enhanced incubation characteristics of the VersaTrek® incubator system. See TREK Diagnostic Systems. The media is available from our diagnostic supply center. Submissions of 3 consecutive draws are recommended. Antimicrobial Susceptibility (MIC) are not included in the fee. See also Foal Diarrhea/Septicemia Kit. *May qualify for New York State contract pricing.	30.00 /sample	Bacteriology	M-F	5-10 days	Whole blood in blood culture medium	40 ml VersaTrek EZ® Draw® REDOX 1® aerobic blood culture bottle	Do NOT Refrigerate
Blood Culture, Anaerobic For the isolation of anaerobic bacteria in blood. Blood culture medium should be inoculated immediately when blood is drawn. See https://ahdc.vet.cornell.edu/docs/Blood_Culture_Technique.pdf for instructions. If the blood culture media used is not the VersaTrek EZ® Draw® REDOX 2® product, turn around time may be delayed due to inability to take full advantage of enhanced incubation characteristics of the VersaTrek® incubator system. See TREK Diagnostic Systems. The media is available from our diagnostic supply center. Submissions of 3 consecutive draws are recommended. Antimicrobial Susceptibility (MIC) are not included in the fee. See also Foal Diarrhea/Septicemia Kit. *May qualify for New York State contract pricing.	30.00 /sample	Bacteriology	M-F	5-10 days	Whole blood	40 ml VersaTrek EZ® Draw® REDOX 2® anaerobic blood culture bottle	Do NOT Refrigerate
Blood Smear Evaluation RBC (morphology, parasites), manual differential cell count, WBC, Plate smear est. Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panelsFor instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear	17.00	Clinical Pathology	M-Sa	1-2 days	1) EDTA blood (> 1/2 full) AND 2) 2 dry (unfixed, unstained) blood smears	1) lavender top tube; 2) frosted-end slides	Refrigerate blood; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment.
Bone Marrow Cytology Report includes – Specimen type and site (provided by submitter), Smear description (overall cellularity, assessment of all hematopoietic lineages, other cellular components and marrow iron by Prussian blue staining), Smear interpretation. Interpretation of bone marrow smears requires a recent CBC (within 24-48 hours). All bone marrow samples without a recent CBC should be accompanied by an EDTA blood sample with 2 blood smears and a CBC should be requested (use 'Other diagnostic tests requested' field to the right of the Clinical Summary field to request a CBC). Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing For instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear	60.00	Clinical Pathology	M - F	1 business day	1) At least 4-10 dry (unfixed, unstained) bone marrow smears	1) Frosted-end slides	Hold at room temperature; DO NOT refrigerate smears; protect from chilling during shipment with insulation.
Bovine Metabolic Panel BHBA, NEFA, UN, Albumin, AST, Cholesterol, Triglycerides Includes NEFA, BHBA, AST, ALB, and BUN; for additional info see Clin Path panels	30.50	Clinical Pathology	M-Sa	1-2 days	2 mL separated serum	plain glass or plastic tube	Refrigerate

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Bronchoalveolar Lavage Count Cytology (smear exam and count: 56.00		Clinical Pathology	M - F	1 business day	1) BAL fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate. Ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect slides from chilling/coolant by insulating separately during shipment.
<p>Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation, respiratory</p> <p>Please also submit 2-3 freshly prepared smears of the BAL fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p>							
Bronchoalveolar Lavage Cytology (smear exam only)	48.00	Clinical Pathology	M - F	1 business day	1) Smears (at least 2-3 unstained smears preferred); and/or 2) BAL fluid in EDTA (no cell counts provided); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only. Ship in insulated container with freezer pack for next day delivery. Hold slides/smears at room temperature; protect from chilling/coolant by insulating separately during shipment
<p>Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation</p> <p>With fluid samples of BAL, please also submit 2-3 freshly prepared smears and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p>							
Bronchodilators	165.50 per hour	Toxicology	M-F	1-3 days	1) 10 mL heparinized whole blood; 2) 25 - 50 mL urine	1) Heparinized tube, no gel clot; 2) sterile, leak-proof container	Refrigerate. Ship in insulated container with ice pack.
<p>(Clenbuterol, Albuterol, Turbutaline)</p> <p>Samples arriving at AHDC between October 27 and December 15 will be forwarded to a referral lab for analysis.</p>							
Brucella Card Agglutination	3.50 /animal	Brucellosis	M-F	1-4 days	1) 2 mL serum; 2) clotted whole blood	1-2) red top blood collection tube	Refrigerate. Ship in insulated container with ice pack.
<p>Non-canine</p> <p>This test will not reliably detect <i>Brucella ovis</i>, <i>Brucella mellitensis</i>, <i>Brucella canis</i>, or <i>Brucella suis</i>. It is intended as a screening test for the detection of antibodies against <i>Brucella abortus</i>.</p>							
Brucella Culture, non-canine	25.00 /sample	Bacteriology	M-F	5-14 days	Affected tissue, lesion or fluid aspirate	Amies transport media (swab) or sealed sterile containers	Refrigerate. Ship in insulated container with ice pack.
<p>For the isolation of <i>Brucella</i> species only.</p> <p>*May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Refer to <i>Brucella canis</i> Culture for canine submissions.</p>							
Brucella Fluorescence Polarization Assay (FPA)	4.00	Bacteriology	M - F	1 - 2 days	100 ul Serum, plasma, whole blood	Red top blood collection tube, serum separator tube	Refrigerate
<p>Designed to meet export, and interstate shipping requirements where requested. For canines, see Canine Brucella in the Bacteriology section</p>							
Brucella Fluorescence Polarization Assay (FPA) Tube Test	4.00	Bacteriology	M - F	1 - 2 days	100 ul Serum, plasma, whole blood	Red top blood collection tube, serum separator tube	Refrigerate
<p>Designed to meet export, and interstate shipping requirements where requested; for canines, see Canine Brucella in the Bacteriology section.</p>							
Brucella Plate Agglutination	25.00 /sample	Brucellosis	M-F	1-6 days	1) 2 mL serum; 2) clotted whole blood	1-2) red top blood collection tube	Refrigerate. Ship in insulated container with ice pack.
<p>Non-canine</p> <p>When required for export or emergencies ONLY</p>							
Brucella Tube Agglutination	6.50 /animal	Brucellosis	M-W	2-7 days	1) 2 mL serum; 2) clotted whole blood	1-2) red top blood collection tube	Refrigerate. Ship in insulated container with ice pack.
<p>Non-canine</p> <p>48-hour incubation; to meet export, bull stud, and interstate shipping requirements; NOT FOR CLINICAL APPLICATIONS. For canines, see "Canine Brucella" in the Bacteriology section</p>							

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Cadmium (Atomic Absorption Spectroscopy - Graphite Furnace)	35.00	Toxicology	M-F	3 days	1) 2g tissue; 2) 100g feed	1) sterile container; 2) any container	1) Refrigerate; 2) none
Calcium Measures total calcium in serum, plasma and fluids. See Eclinpath for more information	11.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Calcium, urine Measures total calcium levels in urine.	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs.
Campylobacter jejuni Culture For the isolation of Campylobacter jejuni. Test is also part of the Enteric Panel 2 and Enteric Panel 3. Antimicrobial susceptibilities are not applicable. Recent data has shown a dramatic shift among the species of Campylobacter involved in small ruminant and camelid abortion cases. Increases have been documented in the incidence of Campylobacter jejuni abortions with research showing that it has now replaced Campylobacter fetus subsp. fetus as the predominant cause of sheep abortions in the United States. We will continue to offer fluorescent antibody (FA) testing for Campylobacter fetus subsp. fetus, however the culture has been discontinued and replaced with Campylobacter jejuni culture.	25.00 /sample	Bacteriology	M-S	3-5 days	1) Fresh feces or intestinal contents; 2) ligated loop of intestine; 3) Placenta (preferred) or fetal liver for small ruminant and camelid abortion cases; 4) Swab of 1 or 2	1-3) Sealed sterile separate containers (sterile preferred); 4) Swab in Amies transport media (w/charcoal preferred)	1-2) Refrigerate
Cantharidin (Gas Chromatography/Mass Spectrometry)	165.50 /sample	Toxicology	M-F	1-3 days	1) urine; 2) stomach contents, vomitus	1-2) sterile, leak-proof container	Refrigerate w/ cool pack
Cerebrospinal Fluid Cytology Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation Specify if the sample is from the atlanto-occipital or lumbosacral region as result interpretation is dependent on site. Do not allow coolant to come in contact with specimen: ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing	58.00	Clinical Pathology	M - F	1 business day	1) CSF in EDTA; 2) For best results also submit freshly made, rapidly air-dried, un-stained smears. 3) Also submit a red top tube if culture anticipated	1) Lavender-top tube; 2) Also submit a red top tube if culture anticipated	1,3) Refrigerate fluid samples; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment.
Chloride See Eclinpath for more information	9.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
Chloride, urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs.
Cholesterol Dogs and cats should be fasted for around 12 hours to avoid cholesterol interference from post-prandial hyperlipidemia. See Eclinpath for more information	13.00 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid;	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
Cholinesterase Measurement of cholinesterase activity in serum or plasma is a quick screening test indicated for animals with a history of possible exposure to organophosphate or carbamate compounds and/or that show clinical signs compatible with exposure. Remember that erythrocytes are rich in cholinesterase, therefore hemolysis invalidates the results.	8.50 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Chromatography Quantitation Special request only, contact Lab at 607-253-3975 or -3974	165.50 per hour	Toxicology	M-F	7 days	1) 10 mL EDTA; 2) 10 mL heparin whole blood; 3) 50 mL urine	Heparinized or EDTA tube, no gel clot; sterile, leak-proof container	1-3) Refrigerate w/ cool pack

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 7 of 30

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Clostridium chauvoei FA For the detection of Clostridium chauvoei by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing .	14.00 /sample	Bacteriology	M-F	1-2 days	1) 2-inch cube of affected muscle; 2) Swab from affected lesions	1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM)	1) Refrigerate; 2) DO NOT refrigerate
Clostridium difficile Toxins (A/B) For the detection of Clostridium difficile toxins A and B by ELISA in enteric samples. *May qualify for New York State contract pricing. For investigational purposes only. The sensitivity of this Clostridium difficile toxin test is maximized by freezing fecal samples that can not be sent chilled to the lab to arrive the next day, since these toxins are heat labile. If you are requesting other testing on feces or colon contents, such as parasitology exams for which the sample should not be frozen, it requires having two separate fecal specimens. The specimen held over a night or weekend can be frozen, labeled as "held frozen" and shipped with extra insulation/freezer packs. A negative toxin test result on a sample that has not been handled in this way should be interpreted with caution as it may not reflect the actual disease status of the animal. Please consider the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions.	33.00 /sample	Bacteriology	M, W, F	2 days	Fresh feces or tied-off loop of bowel (with contents)	Sealed sterile container. Swabs are NOT acceptable.	Refrigerate. Fresh samples must reach the lab within 24 hours of collection. If this is not possible, freeze immediately and ship overnight frozen on dry ice.
Clostridium novyii FA For the detection of Clostridium novyii by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing .	14.00 /sample	Bacteriology	M-F	1-2 days	1) 2-inch cube of affected muscle; 2) Swab from affected lesions	1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM)	1) Refrigerate; 2) DO NOT refrigerate
Clostridium perfringens Enterotoxin For the detection of Clostridium perfringens enterotoxin by ELISA. This ELISA tests for only one of several possible toxins. Please also consider anaerobic culture. The sensitivity of this Clostridium perfringens toxin test is maximized by freezing fecal samples that can not be sent chilled to the lab to arrive the next day, since these toxins are heat labile. If you are requesting other testing on feces or colon contents, such as parasitology exams for which the sample should not be frozen, it requires having two separate fecal specimens. The specimen held over a night or weekend can be frozen, labeled as "held frozen" and shipped with extra insulation/freezer packs. A negative toxin test result on a sample that has not been handled in this way should be interpreted with caution as it may not reflect the actual disease status of the animal. Please consider the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions.	28.00 /sample	Bacteriology	M, W, F	2 days	Fresh feces or tied-off loop of bowel (with contents)	Sealed sterile container. Swabs are NOT acceptable.	Refrigerate. Fresh samples must reach the lab within 24 hours of collection. If this is not possible, freeze immediately and ship overnight frozen on dry ice.
Clostridium septicum FA For the detection of Clostridium septicum by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing .	14.00 /sample	Bacteriology	M-F	1-2 days	1) 2-inch cube of affected muscle; 2) Swab from affected lesions	1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM)	1) Refrigerate; 2) DO NOT refrigerate
Clostridium sordellii FA For the detection of Clostridium sordellii by fluorescent antibody stain. Ship overnight. *May qualify for New York State contract pricing .	14.00 /sample	Bacteriology	M-F	1-2 days	1) 2-inch cube of affected muscle; 2) Swab from affected lesions	1) Sealed sterile container; 2) Anaerobic transport media (BBL™ Port-a-cul™ or Anaerobic Systems ATM)	1) Refrigerate; 2) DO NOT refrigerate
Clotting Factor (07) FVII Activity, Large-Exotic	34.00	Comparative Coagulation		2-3 days	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate
Clotting Factor (09) FIX Activity, Large/Exotic Factor IX coagulant activity	34.00	Comparative Coagulation	M-F	2-3 days	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate
Clotting Factor (10) FX Activity, Large-Exotic The measurement of Factor X coagulant activity in large and exotic animals.	34.00	Comparative Coagulation	M-F	2-3 days	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Clotting Factor (11) FXI Activity, Large-Exotic The measurement of Coagulant Factor XI in large and exotic animals	34.00	Comparative Coagulation	M-F	2-3 days	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate
Clotting Factor (12) FXII Activity, Large-Exotic The measurement of Coagulation Factor XII in large and exotic animals.	34.00	Comparative Coagulation	M-F	2-3 days	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate
Clotting Factor (13) FXIII Deficiency Screening Requires submission of paired sample from same-species control animal	56.00	Comparative Coagulation	Monday - Thursday	1 - 3 days	citrate plasma	plastic or siliconized glass tube in styrofoam box	
Coagulation Panel, Large Animal Incl. aPTT, PT, Fibrinogen Ship Overnight: for additional info on Submission Guidelines, or see Comp Coag web page	42.50	Comparative Coagulation	M-F	same day	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Copper, serum/plasma Contact Lab for specifics for avian and very small animals	25.00 /sample	Toxicology	M-F	3 days	1) 2 mL serum; 2) or transfer 2 mL heparinized plasma into a red top tube	1-2) Red top tube, no gel clot	1-2) Refrigerate. Ship in insulated container with ice pack.
Coronavirus PCR, Alpha The alpha coronavirus PCR employed by the AHDC will detect feline enteric coronavirus, canine enteric coronavirus, ferret or mink coronavirus, and swine coronavirus - AKA transmissible gastroenteritis virus. For species specific testing information please see Coronavirus PCR Guidelines.	36.75	Molecular Diagnostics	M-F as needed	3-5 days	1) EDTA blood, serum (min. 2 mL); 2) tissue (min. 5 g); 3) ascites or body fluid (min. 2 mL); 4) feces (min 5 g)	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice pack.
Corticosteroids, Drug Screen capillary GC/MS Special request only ; contact Lab at 607-253-3975 or -3974. The list of drugs covered in this screen include: Dexamethasone, Betamethasone, Methyl Prednisolone, Prednisolone, Prednisone, Flumethasone, Triamcinolone, and Triamcinolone Acetonide.	331.00 /sample	Toxicology	M-F	3 days	1) 20 mL heparin whole blood; 2) or transfer 5 mL heparin plasma into a red top tube; 3) 50 mL urine	1) Heparinized tube, no gel clot; 2) red top blood collection tube; 3) sterile container	1-3) Refrigerate w/ cool pack
Creatine kinase CK, CPK See Eclinpath for more information	8.50 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Creatinine See Eclinpath for more information	11.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Creatinine, urine	19.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs
Cryptosporidium ELISA ELISA plus Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate. Fecal Qualitative or Quantitative exam will be performed with each ELISA at an additional fee. This ELISA cannot be requested alone. If both Cryptosporidium and Giardia ELISA's are ordered, only one flotation will be performed and charged.	25.00	Parasitology	Mon, Thur	1-3 days	feces	wide mouth leak-proof container	Refrigerate
Cryptosporidium Spp real-time PCR This real-time PCR assay detects the 18S rRNA gene of pathogenic Cryptosporidium species according to the method published by Operario et al. Am J Trop Med Hyg. 2015. This test is intended primarily for research applications at this time. Please contact lab for volume discounts. Ct values at or below 28 are more likely to be associated with active diarrhea. Ct values above 32 are less likely to be associated with diarrhea.	36.75	Molecular Diagnostics	M-F	3-5 days	Feces, intestinal tissue	Sealed vial or container in a secondary container	Refrigerate. Ship in insulated container with ice pack.
Cyanide 1-2) Refrigerate promptly; Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes)	55.25 /sample	Toxicology	M-F	2 days	1) 10 mL EDTA whole blood; 2) 10 mL heparinized whole blood; 3) 50g stomach contents; 4) 100g forage, silage, or other material 3-4) freeze samples promptly, keep samples cold in transit: special request only, contact Lab to make arrangements	Heparinized or EDTA tube; air-tight container	1-2) Refrigerate; 3-4) freeze

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Cytology Smear Exam Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation With fluid aspirates, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing	48.00	Clinical Pathology	M - F	1 business day	1) Smears (at least 2-3 unstained smears preferred); and/or 2) Fluid in EDTA (no cell counts or total protein provided)	1) Frosted-end slides; 2) Lavender-top tube (if submitting fluid)	Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears/slides at room temperature; protect from chilling/coolant by insulating separately during shipment
Dermatology Biopsy Surcharge For Dermatology biopsies (up to 5 sites)	15.00	Anatomic Pathology	M - F	4 - 10 days	0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio.	Wide-mouth, leakproof, unbreakable jar; NOT GLASS	room temperature; protect from freezing
DIC Panel, Large Animal and Exotics Includes aPTT, PT, Fibrinogen, Antithrombin, D-dimer Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page	103.00	Comparative Coagulation	M-F	same day	2 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Dilute Russell's Viper Venom Time Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page	15.50	Comparative Coagulation	M-F	same day	citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Drug Quantitation Special request only, contact Lab	165.50 per hour	Toxicology	M-F	5 days	1) 10 mL heparin plasma transferred to red top; 2) 20 mL heparin whole blood; 3) 50 mL urine	1) red top blood collection tube; 2) Heparinized tube, no gel clot; 3) sterile container	1-3) Refrigerate w/ cool pack
Electrolytes Panel Na, K, Cl For more information, see Chemistry Panel Information	17.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs
Electrolytes Panel, urine Na, K, Cl, Creat For more information, see Chemistry Panel Information.	31.00	Clinical Pathology	M-Sa	1-2 days	2 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs
Electrophoresis, blood	60.00	Clinical Pathology	1x/wk	7-10 days	1) 1 mL separated serum; 2) heparinized plasma for exotics/avian only	plain glass or plastic tube	Refrigerate
Electrophoresis, fluid All species; please specify site of fluid collection	60.00	Clinical Pathology	1x/wk	7-10 days	Body fluids. Please specify site.	plain glass or plastic tube	Refrigerate
Electrophoresis, urine	60.00	Clinical Pathology	1x/wk	7-10 days) 5-7 mL urine	plain glass or plastic tube	Refrigerate
Encephalomyocarditis SN Serum neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation.	16.50 /sample	Virology	as needed	5-9 days	1 ml serum	red top blood collection tube	It is recommended to ship samples with an ice pack.
Encephalomyocarditis Virus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh heart, brain, spleen	Sealable plastic bag or container	Ship in insulated container with ice pack

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Encephalomyocarditis Virus Isolation Virus isolation Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing.	70.00 /animal	Virology	M-F	21-30 days,	s Fresh heart, brain, spleen	Sealable plastic bag or container	Ship in insulated container with ice pack
Enteric Bacteriology Cultures, Panel 1 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], and Salmonella culture. *May qualify for New York State contract pricing. Aerobic culture not tested on calves over 10 days of age. Antimicrobial Susceptibility (MIC) are not included in the testing fee. Please consider the Foal Diarrhea/Septicemia Kit for foal submissions.	62.00 /sample	Bacteriology	M-Sa	3-7 days	1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) Swab of 1-3	1) Sealed, sterile container or ParaPak; 2-3) Sealed sterile container; 4)Swab in Amies transport media	1-3) Refrigerate. Ship in insulated container with ice pack. EXCEPT ParaPak, which is shipped at room temperature and protected from temperature extremes.
Enteric Bacteriology Cultures, Panel 2 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], Salmonella culture and Campylobacter jejuni culture. Antimicrobial Susceptibility (MIC) are not included in the testing fee. Susceptibilities not applicable to Campylobacter jejuni.	87.00 /sample	Bacteriology	M-Sa	3-7 days	1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4)Swab of 1-3	1) Sealed, sterile container or ParaPak; 2-3) Sealed sterile container; 4)Swab in Amies transport media	1-3) Refrigerate. Ship in insulated container with ice pack. EXCEPT ParaPak, which is shipped at room temp and protected from temperature extremes.
Enteric Bacteriology Cultures, Panel 3 Includes aerobic culture [E. coli, Rhodococcus (foals only), Shigella (primates only)], Salmonella culture, Campylobacter jejuni culture, anaerobic culture [Clostridium perfringens (all species) and Clostridium difficile* (foals and primates only)] Antimicrobial Susceptibility (MIC) are not included in the testing fee. Susceptibilities are not applicable to Campylobacter jejuni and Clostridia species. *Culture for Clostridium difficile in other species available by request only. *Culture for Clostridium difficile in other species available by request only. **Anaerobic Transport medias (e.g. BBL™ Port-a-cul™ or Anaerobic Systems ATM) are suitable for most aerobic cultures, including for the aerobes of concern in enteric infections. Despite labelling to the contrary, culture systems containing Amies transport media or Stuart transport media are inappropriate as transport systems for anaerobic culturing. Swabs transported without any transport media are unsuitable for both aerobic and anaerobic culture.	132.00 /test	Bacteriology	M-Sa	3-7 days	1) Anaerobic transport media inoculated with swab of fresh feces or intestinal contents; or 2) Ligated ileum or jejunum; and/or 3) Mesenteric lymph node	1) Anaerobic transport media** (BBL™ Port-a-cul™ or Anaerobic Systems ATM); or 2-3) Sealed sterile container	1) DO NOT Refrigerate anaerobic transport media tubes or vials; 2-3) Tissues are ideally held and shipped frozen with sufficient insulation and freezer packs to arrive next day and very cold; refrigerated temperatures are acceptable only for tissue samples delivered to the lab within 24 hours of collection.
Environmental Samples (for parasites) Please contact Lab: fee depends on parasite of interest, type of sample submitted, and processing required	25.00 See comments for fee	Parasitology	M-F	1-7 days	Please contact Lab before submitting sample to determine if sample is appropriate for testing.		
Ethylene Glycol Special request only, contact Lab	55.00 /sample	Toxicology	M-F	2 days	1) 10 mL heparin whole blood; 2) 10 mL EDTA whole blood; 3) urine; 4) food materials; 5) stomach contents	1,2) Heparinized or EDTA tube, no gel clot3-5) sterile, leak-proof container	1-5) Refrigerate w/ cool pack
Fecal Qualitative A double centrifugation concentration fecal flotation with sugar and/or zinc sulfate solution used as a general test for evaluating feces for parasitic infections The qualitative fecal flotation, using a double centrifugation concentration technique, is a general broad based test for evaluating feces for parasitic infections. It indicates the presence or absence of patent protozoan or worm infections in domestic and wild animals of all species. Samples from individual animals are preferred. Composite samples tell when there is a parasite problem, but not the animals that are particularly affected. In this laboratory, all samples are actively floated (centrifuged) using a sugar solution (1.33 specific gravity). Zinc sulfate solution (1.18 specific gravity) is used also for samples from small animals, animals under six months of age, and when delicate protozoa, such as Giardia, or worms, such as nematode larvae, are of concern. Using these flotation media, protozoan cysts and worm eggs and larvae are recovered for microscopic evaluation. When submitting a sample provide a sufficient sample to make a diagnosis possible, and include a history of the health concerns related to the animal(s) from which the sample(s) are collected. For best results, collect fresh samples, keep them refrigerated, and submit them for examination within seven days of collection. Samples that are preserved in 10% formalin or 70% alcohol can also be submitted, but the preservative may compromise the quality of the sample for parasite detection.	25.00	Parasitology	M-F	1-3 days	10g fresh fecal sample, animal 3 days of age or older	Plastic leakproof container	Freezer Packs. Ship Next Day Delivery

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Fecal Quantitative Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate solution. Primarily used to examine feces of camelids, horses, and ruminants, but other species may be requested. Reported as the number of cysts/ oocysts/ eggs/ larvae per gram. Samples for animals under 3 days of age will not be tested.	25.00	Parasitology	M-F	1-3 days	10g fresh fecal sample, animal 3 days of age or older	wide mouth leak-proof container	Refrigerate
Fecal Wet Mount Was previously called "Direct Fecal Smear" Exam for motile parasites. This test for Cornell CUHA only (must be read within 1 hr of fecal production by host). See Trich. Fecal Exam, Staining for Other Protozoa, or Giardia ELISA	5.50	Parasitology	M-F	1 day	10g fresh fecal sample	wide mouth leak-proof container	Do NOT Refrigerate
Fibrinogen Quantitative measure of clottable fibrinogen (Clauss method) Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page	22.00	Comparative Coagulation	M-F	same day	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Fibrinogen (antigen) Fibrinogen concentration; ELISA method Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page	43.00	Comparative Coagulation	Tu, F	1-3 days	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Fibrinogen (heat-precipitation) Formerly named Fibrinogen (Schalm's) Do not allow coolant to come in contact with specimen. This test is not recommended for small animals due to its inaccuracy and insensitivity. The preferred method for small animals is measurement of clottable fibrinogen (citrate plasma) offered in the Comparative Coagulation lab.	16.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate
Field Necropsy Surcharge This charge will be added to any histopathology request on ante-mortem tissues unless deemed unnecessary by the reviewing pathologist, in which case only a standard histopathology charge will be applied. For necropsied animals it is always recommended to submit a complete set of tissues representing major organ systems and any sites potentially involved based upon clinical history, ante-mortem testing, or post-mortem gross lesions. Tissues will be selected for histopathological examination based upon the clinical history and gross description provided, at the discretion of the pathologist. A clinical history for the individual animal and potentially also for the herd or population from which the animal originates is very important. Also indicate if the animal was euthanized.	56.00	Anatomic Pathology	M-F	4-10 days	0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio.	Wide-mouth, leakproof, unbreakable jar; NOT GLASS	room temperature; protect from freezing
Fluphenazine (LC/MS) (Liquid Chromatography/Mass Spectrometry) Liquid Chromatography/Mass Spectrometry (LC/MS)	165.50 /sample	Toxicology	M-F	1-3 days	1) 10 mL heparine whole blood; or 2) 5 mL heparine plasma transferred to red top tube	Heparinized tube, no gel clot	Refrigerate with cold pack
Formalin-fixed tissue extraction surcharge (Molecular Diagnostics 20.00) Surcharge added to fixed tissues due to the extra labor and materials needed for the nucleic acid extraction procedure.	20.00	Molecular Diagnostics	M-F		Formalin-fixed tissues		Room temp
Fructosamine See Eclinpath for more information	36.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum	red top blood collection tube	Refrigerate. Ship overnight in insulated container with freezer packs
Fungal Culture For the isolation of yeasts, dermatophytes and other filamentous fungi. *May qualify for New York State contract pricing. **Swabs are not encouraged for tissues or body fluids. Keep tissues moist. Ship overnight. Antifungal susceptibilities are not included in the testing fee.	47.00 /sample	Bacteriology	M-F	28+ days	1)Nails, skin scrapings or 10-12 hairs from border of lesion; combings from asymptomatic cats (dermatophytes); 2)Fresh tissue or body fluids (systemic fungi/yeast); 3)Ear (yeast); 4) Swab of lesion, tissue, fluid or ear	1) DRY sterile sealed container (e.g. closed urine cup or envelope); 2) Fluid or tissue in Sterile, sealed container (e.g. red top blood collection tube); 4) Swab in Amies transport media**	1) Not required; 2) Refrigerate
Fungal ID For the identification of pure fungal isolates (filamentous fungi or yeast). Submissions accepted from Laboratories only. Ensure purity before shipment. Follow all biohazard shipping regulations. Provide animal history and suspected identification if possible. Antifungal susceptibilities are not included in the testing fee.	47.00 /isolate	Bacteriology	M-F	28+ days	Fungal isolate (pure culture)	Agar slant or plate (sealed)	Refrigerate

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Fungal KOH Fluorescent Stain For the direct microscopic examination for yeast and fungi. *May qualify for New York State contract pricing . Swabs are not acceptable	12.00 /sample	Bacteriology	M-F	1-2 days	1)Nails, skin scrapings or 10-12 hairs from border of lesion; combings from asymptomatic cats (dermatophytes); 2)Fresh tissue or body fluids (systemic fungi/yeast); 3)Prepared slide	1) DRY sterile sealed container (e.g. closed urine cup or envelope); 2) Sterile sealed container (e.g. red top blood collection tube); 3) Slide mailer	1,3) Not Required; 2) Refrigerate
Fungal Serology Panel For the detection of antibody to Histoplasma, Blastomyces, Coccidioides and Aspergillus (AGID) and Cryptococcus antigen in serum. Tests can be requested individually. See Histoplasma AGID, Blastomyces AGID, Coccidioides AGID, Aspergillus AGID or Cryptococcus antigen.	64.00	Bacteriology	M, W, F	3-5 days	1) Serum; 2) Cerebrospinal fluid (for Cryptococcus antigen only)	1) Red top blood collection tube; 2) Sealed sterile container	1-2) Refrigerate
Fungal Serology: Aspergillus Antibody For the detection of Aspergillus antibody in serum by AGID. Test is also part of the Fungal Serology Panel.	16.00 /sample	Bacteriology	M, W, F	3-5 days	Serum	Red top blood collection tube	Refrigerate. Ship in insulated container with ice pack.
Fungal Serology: Blastomyces Antibody For the detection of Blastomyces antibody in serum by AGID. Test is also part of the Fungal Serology Panel.	16.00 /sample	Bacteriology	M, W, F	3-5 days	Serum	Red top blood collection tube	Refrigerate. Ship in insulated container with ice pack.
Fungal Serology: Coccidioides Antibody For the detection of Coccidioides antibody in serum by AGID. Test is also part of the Fungal Serology Panel.	16.00 /sample	Bacteriology	M, W, F	3-5 days	Serum	Red top blood collection tube	Refrigerate
Fungal Serology: Cryptococcus Antigen For the detection of antigen to Cryptococcus spp. in serum. Test is also part of the Fungal Serology Panel.	22.00 /sample	Bacteriology	M-F	1 day	1) Serum; 2) Cerebrospinal fluid	1) Red top blood collection tube; 2) Sealed sterile container	1-2) Refrigerate
Fungal Serology: Histoplasma Antibody For the detection of Histoplasma antibody in serum by AGID. Test is also part of the Fungal Serology Panel.	16.00 /sample	Bacteriology	M, W, F	3-5 days	Serum	Red top blood collection tube	Refrigerate
Fungal Susceptibility This test includes Amphotericin B, 5-Flucytosine, Anidulafungin, Caspofungin, Micafungin, Fluconazole, Itraconazole, Posaconazole, Voriconazole *May qualify for New York State contract pricing. **Swabs are not encouraged for tissues or body fluids. Susceptibilities will NOT be performed on dermatophytes or other slow growing fungi. Fungal culture MUST be performed to recover an isolate before testing can be done. ***Lag time may be longer if isolation is required. From time of isolation: 5-6 days for yeast, 12-14 days for filamentous fungi.	68.00 /isolate	Bacteriology	M-TW	6 days for pur	1) Fresh tissue or body fluids (for filamentous fungi and yeasts to assess systemic infection); 2) Fungal isolate (pure culture) – yeast or fast growing filamentous fungi ONLY	1) Sealed sterile container (preferred) or *Amies transport media (swab); 2) Agar slant or plate (sealed)	1) Refrigerate; 2) Not required
Gamma Glutamyl Transferase (GGT) See Eclinpath for more information	9.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	Red top (non-anticoagulant) or green top (heparin) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
Gamma Glutamyl Transferase (GGT), urine	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs.
Giardia ELISA ELISA plus Centrifugation Concentration Flotation with Sugar and/or Zinc Sulfate. A Fecal Qualitative or Quantitative exam will be performed with each ELISA at an additional fee. This ELISA cannot be requested alone. If both Cryptosporidium and Giardia ELISA's are ordered, only one flotation will be performed and charged.	25.00	Parasitology	Mon, Thur	1-3 days	feces	wide mouth leak-proof	Refrigerate

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Glucose Glucose levels in blood and fluids Delayed serum separation or submission of whole blood (cells on serum or plasma) results in a falsely decreased (uninterpretable) glucose. Serum should also be removed promptly from serum separator tubes and placed into a new red top tube. See Eclinpath for more information	11.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum, EDTA plasma, or heparinized plasma; 2) fluid	Red top (non-anticoagulant), green top (heparin), or purple top (EDTA) tube	Refrigerate. Ship overnight in insulated container with freezer packs.
Glucose, urine	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	Red top (non-anticoagulant) tube or plastic leakproof container	Refrigerate. Ship overnight in insulated container with freezer packs
Glutamate Dehydrogenase, blood	8.50	Clinical Pathology			1) 1 mL separated serum or heparinized plasma; 2) fluid		
Gram Stain For the direct microscopic examination for bacteria. *May qualify for New York State contract pricing. Please consider using the Foal Diarrhea/Septicemia Kit and Adult Equine Diarrhea Kit for equine submissions. Specific bacteria cannot be definitively identified by this test.	8.50 /sample	Bacteriology	M-F	1-2 days	1) Affected tissue, lesion or fluid; 2) Prepared slide	1) Affected tissue, lesion or fluid in sealed sterile container; 2) Slide mailer	1) Refrigerate; 2) Not required
Hemoglobin (HGB) Do not allow coolant to come in contact with specimen; ship Overnight	7.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate
Hemogram (CBC), Automated (Mammalian) WBC, RBC, HgB, HCT, MCV, MCH, MCHC, RDW, Plate Count, MPV, Automated Differential Cell Count Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panelsFor instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear	18.50	Clinical Pathology	M-Sa	1-2 days	1) EDTA blood (> 1/2 full) AND 2) 2 dry (unfixed, unstained) blood smears	1) lavender top tube; 2) frosted-end slides	1) Refrigerate blood; ship in insulated container with freezer pack for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment.
Hemogram (CBC), Partial WBC, RBC, HgB, Platelet Count, PCV, MPV, RDW, MCH, MCHC, MCV Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Hematology Testing.	11.50	Clinical Pathology	M-Sa	1-2 days	1) EDTA blood (> 1/2 full) AND 2 dry, unfixed, unstained blood smears	1) lavender top tube; 2) frosted-end slides	Refrigerate
Hemogram, Large Animal WBC, RBC, HgB, HCT, MCV, MCH, MCHC, RDW, Plate Count, MPV, Morphology, Manual Differential Cell Count, Plate Smear est, Plasma Appearance, TP Protect slides from chilling/coolant during storage/shipment.For instructions on making blood smears see https://ahdc.vet.cornell.edu/Sects/ClinPath/sample/test/hema.cfm#Bloodsmear	41.00	Clinical Pathology	M - Sa	1 - 2 days	1) EDTA whole blood AND 2) 2 unstained, air-dried blood smears	1) lavender top tube; 2) frosted-end slides	Refrigerate blood; ship in insulated container with freezer packs for next day delivery. Hold slides at room temperature; protect from chilling/coolant by insulating separately during shipment
Heparin (Low Molecular Weight) Anti-Xa activity of low molecular weight heparin Ship Overnight; for determination of peak anti-Xa values, collect sample 2-3 hrs (cat) or 3-4 hrs (dogs) after heparin dose. For additional info on Submission Guidelines, or see Comp Coag web page. Also see Heparin Assay FAQs.	39.25	Comparative Coagulation	M-F	same day	citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Heparin (Unfractionated) Anti-Xa activity of unfractionated heparin Ship overnight; specify UFH or LMWH; for additional info on Submission Guidelines, or see Comp Coag web page. Also see Heparin Assay FAQs. Peak and trough values: samples for determination of Peak anti-Xa activity should be collected at 3 to 4 hours post treatment (dogs) or 2 to 3 hours post treatment (cats). Samples for Trough levels should be collected just before the next scheduled dose.	39.25	Comparative Coagulation	M-F	same day	0.5 to 1.0 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Histopathology	66.50 for 1-2 sites; Field necropsy/veterinarian-performed necropsy (necropsy in a bottle) \$122.50; Dermatology biopsies \$81.50; companion animal liver biopsies \$111.50. Special/augmented histo requests \$20.50 for each additional site beyond 1-2 tissues. If STAT, call the lab in advance. See notes below regarding what is included in each of these categories of histology requests."	Anatomic Pathology	M-F	4-10 days	0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio.	Wide-mouth, leakproof, unbreakable jar; NOT GLASS	room temperature; protect from freezing
<p>Microscopic examination of formalin fixed, paraffin embedded tissue slides with hematoxylin and eosin staining. Additional special stains, including up to two immuno-histochemistry (IHC) stains, will be added if deemed necessary by the pathologist. Additional IHC stains may be requested for an additional fee.</p> <p>The 1-2 site fee is intended for surgical biopsies and is the baseline charge for the special biopsy categories below. Surcharges for these will be added to the surgical biopsy fee based on the following criteria: Dermatology biopsies: Up to five sites or tissue punches. Please use the Dermatohistopathology form to request this service. For more information, please see Dermatopathology Samples. Liver Biopsy: All canine and feline liver biopsies submitted through the surgical pathology service will receive a surgical biopsy charge plus an additional \$45.00 surcharge for a liver special stain panel. This additional charge will cover a series of special stains and review by our liver pathology specialist since the H&E stain alone may be insufficient to provide proper hepatic analysis. Upon request, a Digital Copper Quantification (DCQ) and/or consultation with a clinical specialist in hepatic disease can be added for additional fees. Please use the Small Animal Liver Biopsy form to request these services. Due to the need for these special stains the processing of reports for these specimens may not be available for 7-10 business days. Special/Augmented biopsies: Intended for complex submissions of entire organs, lobes, systems, tumor beds, limbs, jaws, etc. This includes research/insurance/legal cases for which every single tissue submitted must be described histologically and the selection of the tissues is not at the discretion of the pathologist. It may also include complicated submissions of entire organs or tumor beds or lymphatic samples that need exact tracking of site of collection for multiple samples of the same tissue type. Please call 607-253-3319 in advance to discuss research histology needs or for price quotes related to these types of submissions. Field Necropsy: For necropsied animals it is always recommended to submit a complete set of tissues representing major organ systems and any sites potentially involved based upon clinical history, ante-mortem testing, or post-mortem gross lesions. Tissues will be selected for histopathological examination based upon the clinical history and gross description provided, at the discretion of the pathologist. A clinical history for the individual animal and potentially also for the herd or population from which the animal originates is very important. Also indicate if the animal was euthanized. Please contact the lab to arrange for STAT testing. Three tissues per case will be reviewed as a STAT unless prior arrangements are made. Call 607-253-3319 or email pathology service @cornell.edu. *May qualify for New York State contract pricing</p>							
Histopathology (Each additional site)	20.50 each additional site	Anatomic Pathology	M-F	4-10 days	0.5 - 1 cm thick section of tissue fixed in 10% neutral buffered formalin, 10:1 formalin to tissue ratio	Wide-mouth, unbreakable jar containing 10% neutral buffered formalin; NOT GLASS	
<p>Additional charge added for each site evaluated beyond 2 that are part of a standard biopsy. These additional fees will be applied to submissions of multiple masses from a single organ that require separate slides to track specific anatomic locations and monitor margins. They will also apply to submissions for research and legal purposes, as well as general purposes, for which every submitted tissue must be examined. Please call the lab to arrange STAT testing. Three tissues per case as a STAT unless prior arrangements are made. Please call 607-253-3319 to arrange STAT histopathology.</p>							
Histopathology, Slide Review (1-2 slides)	55.00	Anatomic Pathology	M-F	3 days	1 to 2 Slides	Slide mailer	
<p>1-2 slides Contact Path Dept (607-253-3312)</p>							
Histopathology, Slide Review (3+ slides)		Anatomic Pathology	M-F	3 days	3 or more slides	Slide mailer	
<p>3 or more slides Contact Path Dept (607-253-3312)</p>							
IHC w/o Histopathology	35.00	Anatomic Pathology	M-F	2-7 days	formalin fixed tissue (paraffin-embedded), or tissue mounted on glass slides		room temperature; protect from freezing

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
 VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Influenza Virus Matrix PCR This PCR assay will detect most if not all influenza virus types currently circulating in most species, including 2009 H1N1 and the 2015 Canine H3N8. This assay does not specifically identify the influenza H and N types. That testing can potentially be requested on positive samples if unusual outbreaks or cross-species transmission is suspected. Submit swabs in sterile, sealed vials with several drops of saline added, to prevent desiccation. Bacterial transport media are not appropriate for PCR assays. If culture is desired, submit a separate swab in an acceptable bacterial transport media. Cotton, plastic, wood-handled, and Dacron and other synthetic swabs are all acceptable for PCR-based assays. Calcium alginate swabs should be avoided. All canine and feline samples testing positive for influenza A matrix will be further tested to determine subtype. Subtyping will be done automatically and at no additional charge. Results will be available within 3-5 days following release of the original result on an addended report. *May qualify for New York State contract pricing. This service is performed pursuant to an agreement with Roche Molecular Systems.	36.75	Molecular Diagnostics	M-F as needed	3-5 days	1) tracheal, choanal, oropharyngeal, cloacal (duck) 2) tissue (min. 5 g); 3) allantoic fluid (min. 2 mL)	unbreakable, leak-proof container	Refrigerate
Ionized Calcium Test run on serum only For special sample collection instructions, see Appendix A	18.50	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum ONLY (see Appendix A)	non-anticoagulant tube (plain red top)	Refrigerate
Ionophore Quantitation Monensin, Lasalocide, Salinomycin Liquid Chromatography/Mass Spectrometry (LC/MS)	165.50 per hour	Toxicology	M-F	1-3 days	1) feed; 2) 10 mL heparin whole blood	1) leak-proof zip-lock bag; 2) green top blood collection tube	Refrigerate w/ cool pack
Ionophore Screen Monensin, Lasalocide, Salinomycin Liquid Chromatography/Mass Spectrometry (LC/MS)	330.75 /sample	Toxicology	M-F	1-3 days	feed samples	leak-proof zip-lock bag	Refrigerate w/ cool pack
Iron	8.50 /sample	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma;	plain glass or plastic tube	Refrigerate
Iron Panel Iron (FE), Total Iron Binding Capacity, % Saturation for additional info on Clin Path panels	20.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Knott's Heartworm Used to assess microfilarial burden or to differentiate <i>Dirofilaria immitis</i> and <i>Dipetalonema reconditum</i> microfilariae. If an Occult heartworm antigen ELISA test is also desired, the laboratory can separate plasma from the EDTA or heparin whole blood (if sufficient volume) or a serum sample can be submitted. Please clearly indicate each test as a unique test request.	16.50	Parasitology	M-F	1-3 days	1 mL EDTA or heparin whole blood	sterile blood tube	Refrigerate
Large Animal Chemistry Panel Equ and ruminant: Na, K, Cl, Bicarb, Anion Gap, BUN, Creat, Ca, P, Mg, TP, Alb, Glob, A/G ratio, Gluc, SDH, AST, GLDH, GGT, Tot/Dir/Ind Bilirubin, CK, Iron, TIBC, % Saturation For more information, see Chemistry Panel Information.	48.00	Clinical Pathology	M-Sa	1-2 days	2 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Large Animal Renal Panel Equ and ruminant: Na, K, Cl, Bicarb, Anion gap, Urea, Creat, Alb, Ca, P	28.00	Clinical Pathology	M-Sa	4 hours	1 mL separated serum	plain glass or plastic tube	Refrigerate
Lawsonia intracellularis qPCR This qPCR assay detects the aspartate ammonia lyase (aspA) gene of <i>Lawsonia intracellularis</i> , the causative agent of proliferative enteropathy (also known as Ileitis), in any host species. Due to the intracellular growth of this bacterium, it cannot be cultured for diagnostic purposes. Based on experimental studies in foals, detection of fecal shedding by PCR is expected to occur at 12-18 days after infection, which is approximately the same time frame as onset of clinical signs; shedding may last for 7-21 days (Pusterla et al. J Vet Intern Med 2010). In pigs, shedding starts one week after infection (just after the onset of clinical signs) and may last intermittently for up to 12 weeks; pigs given the modified-live vaccine may be positive by PCR for up to 9 weeks (Guedes et al. Vet Microbiol 2003). This service is performed pursuant to an agreement with Roche Molecular Systems.	36.75	Molecular Diagnostics	M-F as needed	3 - 5 days	Feces, illium, or tonsillar tissue	Leak-proof container	Refrigerate. Ship in insulated container with ice packs
Lead, blood Lead, blood Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes). For avian and exotics, contact Laboratory	35.00 /sample	Toxicology	M-F	2 days	1) EDTA whole blood (min .5 ml); 2) heparin whole blood (min .5 ml)	Heparinized or EDTA tube	1-2) Refrigerate. Ship in insulated container with ice pack.
Lead, liquids/fluids	30.00 /sample	Toxicology	M-F	1-2 days	1) urine; 2) water; 3) milk	1-3) sterile container	1-3) Refrigerate w/ cool pack

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Lead, tissue/feed/other (Atomic Absorption Spectroscopy - Graphite Furnace)	45.00	Toxicology	M-F	3 days	1) 2g tissue; 2) 100g feed	1) sterile container; 2) any container	1) Refrigerate; 2) none
Leptospira MAT, 5 standard serovars	16.00 /5 serovars	Bacteriology	M, W, F	2-3 days	2.0 ml serum	Red top blood collection tube	Refrigerate. Ship overnight in an insulated container with ice packs
For the detection of antibodies to Leptospira interrogans serovars Pomona, Hardjo, Icterohemorrhagiae/copenhageni, Grippotyphosa, and Canicola by serum microagglutination. Please indicate any previous testing at the AHDC for accurate test result interpretations. Testing paired samples at an interval between 2-6 weeks between blood draws has more diagnostic significance than relying on the results from a single sample. See also Leptospira MAT, 7 serovars for additional serovar options that may be relevant. The first 5 serovars may qualify for New York State contract pricing. Refer to Leptospira Microagglutination Testing at the AHDC for more information.							
Leptospira MAT, 7 serovars	30.00 /7 serovars	Bacteriology	M, W, F	2-3 days	2.0 ml serum	Red top blood collection tube	Refrigerate. Ship overnight in an insulated container with ice packs
For the detection of antibodies to Leptospira interrogans serovars Pomona, Hardjo, Icterohemorrhagiae/copenhageni, Grippotyphosa, Canicola, Autumnalis and Bratislava by serum microagglutination. This test adds Leptospira interrogans serovars Bratislava and Autumnalis to the Leptospira MAT, standard 5 serovars. These two serovars are much less likely to be found in the United States, however, the AHDC has identified clinical samples with strongly elevated titers to one or both of these serovars in the absence of strong reactions to the other five. Testing for these two serovars may therefore be of diagnostic value. Please indicate any previous testing at the AHDC for accurate test result interpretations. Testing paired samples at an interval between 2-6 weeks between blood draws has more diagnostic significance than relying on the results from a single sample. Refer to [Leptospira Microagglutination Testing							
Leptospira real-time PCR	36.75 per specimen tested	Molecular Diagnostics	M-F	3-5 days	1) Urine; or 2) Blood (EDTA whole blood); or 3) Urine and blood (EDTA whole blood); or 4) Kidney and/or liver; or 5) Fetal lung, kidney, liver, stomach or abomasal contents, placenta; or 6) Ocular fluids or 7) Bacterial culture. Refer to Testing Strategy information below.	Leak-proof container	Refrigerate. Ship in insulated container with freezer pack
This assay targets the lipL32 gene encoding the major surface antigen of pathogenic Leptospira species.							
Lipase	24.00 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate
Listeria Culture	22.00 /sample	Bacteriology	M-F	5-7 days	1) Encephalitic: brain stem, cerebellum; 2) Visceral: liver, spleen, kidney; 3) Swab of 1 pr 2; 4) Silage	1-2) Sealed separate sterile containers; 3) Amies transport media (swab); 4) Sealed sterile container	1-3) Refrigerate
For the culture of Listeria monocytogenes ONLY. Neurologic listeriosis is not typically associated with a bacteremia. CNS infection is believed to and has been shown to result from localized entry of the bacteria via defects in the oral mucosa, with initial entry into the CNS via the trigeminal nerve. A systemic or bacteremic phase is not usually associated with this process. CSF or brain would be required to identify neuroinfection in most cases. Blood can be a good sample in monogastrics and neonates with septicemia. Consider submission of inoculated Blood Culture media and requesting "Blood culture, aerobic" for testing a clinically ill live animal. Placenta and fetal tissues are appropriate specimens in abortions cases. Our laboratory does not provide food safety testing for listeria on milk or other foods. Samples submitted for testing should be related to clinical investigations. Antimicrobial susceptibilities are not included in the testing fee. Test is also part of Ruminant CNS Panel. Rabies testing should also be submitted to the NYS DOH Rabies laboratory for neurologic submissions. Preliminary results will be provided one day after sample processing begins.							
Magnesium	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate
Magnesium, urine	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate
Mass Spectroscopy Confirmation	165.50 /sample	Toxicology	M-F	5 days	1) 10 mL heparin whole blood; 2) 20 - 50 mL urine; 3) tissue; 4) plant materials; 5) stomach contents	1) Heparinized tube; 2-5) sterile, clean, pesticide-free container	1-2) Refrigerate w/ cool pack; 3-5) freeze tissue
Freezer pack whole blood: SPECIAL REQUEST - PLEASE CONTACT LAB.							
Mass Spectroscopy Screen	331.00 per sample	Toxicology	M-F	7 days	1) 10 mL heparin whole blood; 2) 20 - 50 mL urine; 3) tissue; 4) plant materials; 5) stomach contents	1) Heparinized tube; 2-5) sterile, clean, pesticide-free container	1-2) Refrigerate w/ cool pack; 3-5) freeze tissue
Mercury	55.00	Toxicology	M-F	3 days	1) 2g tissue; 2) 100g feed; 3) 2 mL heparin whole blood	1) sterile container; 2) any container; 3) Heparinized or EDTA tube	1, 3) Refrigerate; 2) none
(Flameless Atomic Absorption) Cannot analyze in plasma or serum - Need whole blood. Do not use heparin separator tubes (gel tubes).							
Microscopic Feed Analysis	40.00 /hr	Toxicology	M-F	3 weeks	100g feed	any container	None
Mineral Panel (Tissue, Feed)	47.00	Toxicology	1x/wk	3-7 days	1) Tissue; 2) Feed. Do not submit serum or plasma.	1) sterile container; 2) any container	1) Refrigerate; 2) none
Full screen includes Ca, P, Mg, Na, K, Cu, Fe, Mn, Zn, and Mo analyzed by ICP-AES. For biopsy samples and other submissions <0.500 g, only Cu, Fe, Zn, and Mo will be provided due to limitations of small sample size. See Description and Minimum Sample Volume for special note on biopsy or other very small samples.							

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 17 of 30

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Mineral Panel, urine Urine mineral panel includes Magnesium, Phosphorous, Creatinine and Calcium For additional info on Clin Path panels	33.00	Clinical Pathology	M-Sa	1-2 days	2 mL urine	plain glass or plastic tube	Refrigerate
Mineral/Electrolytes Panel Na, K, Cl, Bicarb, Anion Gap, Ca, P, Mg For additional info on Clin Path panels	21.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Mineral/Electrolytes Panel, urine Na, K, Cl, Ca, P, Mg For additional info on Clin Path panels	39.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate
Mycoplasma Culture For the culture of Mycoplasma species. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Test is also part of the Reproductive Panel 4 and Respiratory Panel 6.	27.00 /sample	Bacteriology	M-F	5-10 days	1) Uterine, cervical or vaginal mucus; 2) Semen; 3) Tracheal washing or lung tissue; 4) Fluid (joint, pericardial, urine, etc.); 5) Eye; 6) Ear; 7) Aseptically collected swab of 1-6	1-6) Tissue or fluid in sealed sterile container 7) Swab in Amies transport media WITHOUT charcoal or other suitable mycoplasma culture transport media	1-6) Refrigerate
Necropsy After Hours Fee An after-hours fee of \$50 will be applied when bodies are dropped off after 6pm during the week or between 3pm Saturday and 7:30am Monday morning. Additional STAT fees may apply if a necropsy is required at the time of drop off.	50.00	Anatomic Pathology					
Necropsy, greater than 500 lbs Includes large hoofstock, large carnivores \$100 Stat fee will be charged for nights, weekends or holidays. Stat fees will be waived for zoonotic disease suspects or herd health outbreaks. Please call 607-253-3319 to arrange. For shipping instructions, please see Shipping NecropsySpecimens. To ship by UPS or FEDEX, package must weigh less than 150 lbs. They require notification prior to scheduled pickups for all packages exceeding 75 lbs.	410.00 in state; \$550 out of state. Additional STAT fee may apply. Contact lab for more information	Anatomic Pathology	M-F	2-12 days	carcass		
Nematode Larval Culture Requires 14-day lag time from submission; Fecal Quantitative test done first to determine need	55.00	Parasitology	M-F	14+ days	15g fresh fecal sample	wide mouth leak-proof	Do not refrigerate sample
Nitrate/Nitrite, liquids/fluids	24.00 /sample	Toxicology	M-F	2 days	1) water; 2) serum; or 3) transfer heparin or EDTA plasma to a red top tube	1) sterile leak-proof container; 2-3) red top blood collection tube (no gel separator tubes, please)	1-3) Refrigerate w/ cool pack
Nitrate/Nitrite, solids	33.00 /sample	Toxicology	M-F	2 days	1) stomach or rumen contents; 2) feeds	1) sterile leak-proof container; 2) leak-proof zip-lock bag	1-2) Freeze
Non-Esterified Fatty Acid (NEFA) NEFA for additional info on Clin Path panels	13.00	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum; 2) EDTA plasma	1) red top blood collection tube; 2) purple top collection tube	Refrigerate
Osmolality Do not allow coolant to come in contact with specimen	19.50 /sample	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid; 3) 2 mL urine; 4) milk replacer	plain glass or plastic tube	Refrigerate
Packed Cell Volume PCV or HCT (Hematocrit) Do not allow coolant to come in contact with specimen; ship Overnight	7.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Parasite Identification	28.00	Parasitology	M-F	1-7 days	whole parasites including worms or arthropods; punch biopsy, skin or conjunctival snips, fresh tissue for worm identification; impression smears; stained tissue section slides; environmental samples	plastic leak proof container, such as blood tube, appropriate for sample. Submit fresh, in 70% ETOH or 10% formalin, depending upon the sample.	Refrigerate if not fixed
<p>Endo- and ecto- parasites All ticks submitted for PCR (Lyme, Anaplasma, Ehrlichia) must be identified prior to testing. A biopsy is examined directly or undergoes pepsin digestion to recover nematodes. Gastrointestinal contents and other tissues such as lung are examined and processed as needed to collect and identify parasites present. Intestinal smears are made from fresh, non-preserved intestinal sections and are used to identify non-patent coccidian infections postmortem. Other samples that can be submitted for parasite identification include: Biopsies from tissues that may contain parasites; Necropsy tissues, such as sections of organs and intestines with contents; Fluids such as tracheal and cloacal washes, duodenal aspirates, urine, blood; Histological sections on slides; Free living worms or arthropods found on animals, in residences, in feed samples and environmental samples such as soil and composted material.</p>							
Pericardial Fluid Cytology	50.00	Clinical Pathology	M - F	1 business day	1) Pericardial fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment.
<p>Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With pericardial fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p>							
Peritoneal Fluid Cytology	50.00	Clinical Pathology	M - F	1 business day	1) Peritoneal fluid in EDTA; 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only, DO NOT refrigerate smears
<p>Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With peritoneal fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p>							
Phosphate (Phosphorus)	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate
Phosphate (Phosphorus), urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate
Platelet Aggregation	97.00	Comparative Coagulation	---	---	requires special arrangement with lab		
<p>Combination of individual tests Special arrangement - contact laboratory to schedule testing</p>							
Platelet Count	18.00	Clinical Pathology	M-Sa	1-2 days	1) EDTA blood (> 1/2 full); 2) 2 dry (unfixed, unstained) blood smears	1) lavender top tube; 2) frosted-end slides	Refrigerate
<p>Do not allow coolant to come in contact with specimen or slides; ship Overnight</p>							
Platelet Function	52.00	Comparative Coagulation		2-3 days	requires referral appointment		
<p>Combination of individual tests Special arrangement - contact laboratory to schedule testing</p>							
Pleural Fluid Cytology	50.00	Clinical Pathology	M - F	1 business day	1) Pleural fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment.
<p>Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Smear type, Smear description, Smear Interpretation With pleural fluid samples, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing</p>							

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
 VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Porcine Adenovirus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh lung, intestine, kidney, brain	Sealable plastic bag or container	Ship in insulated container with ice pack
Porcine Hemagglutination Encephalitis Virus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh brain, spinal cord	Sealable plastic bag or container	Ship in insulated container with ice pack
Porcine Parvovirus FA Fluorescent antibody	25.00	Virology	M - F	2-4 days	Fresh fetal lung, intestine, lymph nodes, spleen	Sealable plastic bag or container	Ship in insulated container with ice pack
Porcine Reproductive Respiratory Syndrome FA Fluorescent antibody	25.00	Virology	M - F	2 - 4 days		Sealable plastic bag or container	Ship in insulated container with ice pack
Porcine Rotavirus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh jejunum, ileum	Sealable plastic bag or container	Ship in insulated container with ice pack
Potassium	8.50 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate
Potassium, urine	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate
PreAnesthesia Panel, Large Animal for additional info on Chemistry Panel Information	30.00 /sample	Clinical Pathology	M-Sa	1-2 days	2 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Prothrombin Consumption Index Ship Overnight; tested by special arrangement: contact laboratory for instructions	40.00	Comparative Coagulation	Th-F	1-2 days	paired citrate plasma and serum samples	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Prothrombin Time (PT) Ship Overnight; for additional info on Submission Guidelines, or see Comp Coag web page	14.50	Comparative Coagulation	M-F	same day	1 mL separated citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
Pseudorabies Virus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh brain, spleen, lymph node	Sealable plastic bag or container	Ship in insulated container with ice pack
Pseudorabies Virus Isolation Virus Isolation Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing.	70.00 /animal	Virology	M-F	21-30 days, s	Fresh brain, spleen, lymph node	Sealable plastic bag or container	Ship in insulated container with ice pack
Rabies Brain Extraction This procedure is intended for removal of the brain from the skull for further rabies testing and other diagnostics. A Rabies History Form is required for all rabies suspect cases. Submitting the head or carcass to the Anatomic Pathology service for brain extraction in rabies suspect cases will be followed by shipment of brain tissue to the NYS Dept of Health Lab for Rabies testing. Consideration for field collection of brain specimens and direct shipment to the NYSDOH by attending veterinarians should be considered in case involving human exposure, especially when weekends and holidays will result in further shipping delays. See these links for further information on collection and submission of appropriate specimens for direct submission.	51.00	Anatomic Pathology	M-Sa	24 hrs, follow head or whole body		leak-proof container	Refrigerate, do not freeze
Red Blood Cell (RBC) Count, Total Do not allow coolant to come in contact with specimen; Ship Overnight.	7.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate
Red Blood Cell (RBC), Parasites This test is performed using Wright's (Giemsa) Stain. Do not allow coolant to come in contact with specimen; ship Overnight	12.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
 VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Ref: Aldosterone Assay to quantitate aldosterone levels in EDTA plasma or serum. Can test a single sample for baseline values or can provide pre and post ACTH samples as ACT stimulation test. For All Species: Pre or post samples can be serum, plasma or one of each. Send via overnight courier	48.00 (+ referral fee; see comment)	Referral	Mon and Thurs	4-7 Business	1mL EDTA plasma or serum; serum or EDTA plasma should be separated from clot or cells by centrifugation, respectively, within 30 minutes, and transferred to plain tube with no additives.	Red top, Purple top, or Tiger top blood collection tube	Refrigerate. Ship for next day delivery in insulated container with freezer pack(s)
Ref: Aspergillus Panel This panel includes Aspergillus Antibody Elisa, Electrophoresis to Quantitate host plasma proteins, and galactomannan assay for a dominant Aspergillus antigen. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Testing performed on Avians and Mammals. Sample should be 5 days old or less at time of testing. Do not collect and ship to AHDC prior to 3 day weekend or other conditions which will not allow delivery to referral lab for testing by 5 days post collection. For test interpretations, please see Testing for Avian Aspergillus	102.00	Referral	M - F		0.5 ml heparinized plasma	Green top blood collection tube	Refrigerate, ship sample over night in insulated container with freezer packs.
Ref: Babesia Gibsoni Giemsa Smr slides, Exp Evaluation of blood smears for hemoparasites/ hemotropic bacteria. Also known as Hemoparasite exam – Wright Giemsa Stain Submitter must make blood smear slides. Do not submit EDTA whole blood. Testing can be performed on the following: Amphibian, Avian, Bovine, Canine, Caprine, Cervid, Equine, Exotic, Feline, Ovine, Porcine, Reptile, and Rodents. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	19.00 (+ referral fee; see comment)	Referral	Monday-Friday	2 days	1-2 Unstained, air-dried blood smear slides made from EDTA Whole Blood	slide mailer	Hold at room temperature. Ship at room temperature. Protect from cold temperatures with insulation.
Ref: Babesia PCR This is a polymerase chain reaction assay which amplifies and detects target nucleic acid sequences of the Babesia genus. This laboratory performs nucleic acid sequencing for all positive samples, verifying detection and providing identification to the species and occasionally strain level. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. This test can be performed on all species.	60.00 (+ referral fee; see comment)	Referral	Tues; Thurs	5-7 days	2ml EDTA Whole Blood	Purple Top Collection Container	Refrigerate. Ship in insulated container with freezer packs to arrive next day.
Ref: Bartonella PCR This is a polymerase chain reaction assay which amplifies and detects target nucleic acid sequences of the Bartonella genus. This laboratory performs nucleic acid sequencing for all positive samples, verifying detection and providing identification to the species and occasionally strain level. Sample required is EDTA whole blood. This test can be performed on all species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	60.00 (+ referral fee; see comment)	Referral	Tues, Thurs	5-7 days	2 mL EDTA whole blood	Purple Top blood Collection Container	Refrigerate, send via overnight delivery; insulated container with freezer pack(s)
Ref: Botulism Real-time PCR assay for the detection of Clostridium botulinum types A, B and C toxin genes in vegetative cells after a culture enrichment step ?Finding C. botulinum organisms or spores in a fecal or GI sample of a horse or cow, or in a feed sample, is strongly supportive of a diagnosis of botulism in animals exhibiting compatible clinical signs. Please see http://www.vet.upenn.edu/docs/default-source/new-bolton-center-hospital/NBC-Diagnostic-Labs/instructions-for-submitting-samples-9-1-16.pdf?sfvrsn=6 for more information.	0.00 Cost cannot be estimated prior to consultation with laboratory (+ referral fee; see comment)	Referral			Stomach/rumen contents, GI contents, feces, spoiled forages, parts of dead animals found in suspect feed, and soil from underneath affected areas.	Samples should be sent in whirlpack bags or other suitable small leak proof container. Put each intestinal sample (contents only, please do not send whole intestines or ligated intestines) in a separate Whirlpak bag and clearly label with ID with a waterproof marker such as a Sharpie.	Refrigerate for immediate shipment or store frozen; Ship for next day delivery in insulated container with freezer pack(s)
Ref: Brach sequencing PCR This test sequences the Nox gene from Brachyspira cultures for speciation. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	60.00 (+ referral fee; see comment)	Referral	as needed	3-5 days	identification from culture	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Ref: Brachyspira culture Culture for Brachyspira spp. bacteria. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	30.00 (+ referral fee; see comment)	Referral			rectal swab in transport media, feces, colon, or cecum	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs
Ref: Brucella abortus/suis Fluorescent Polarization Assay This is an antibody detection assay that is used primarily for regulatory diagnostic investigations or for export purposes. It detects antibodies against Brucella abortus or Brucella suis. Test can be performed on large animals. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	33.60	Referral	Mon, Thurs	5-7 days	2mL Serum	Red top blood collection tube.	Refrigerate; ship via overnight delivery in insulated container with freezer packs.
Ref: Cholinesterase(brain) This assay tests for enzyme activity in a brain sample. Any factor that interferes with enzyme activity will affect the outcome of the results. Specimen should be as fresh as possible. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	48.00 (+ referral fee; see comment)	Referral	M-F	3-5 days	Brain Tissue/Retina. Split brain into two equal halves sagittally to get a representative hemisphere. Retina-whole eye.	Plastic bag	Keep refrigerated at all times. Ship overnight with ice pack.
Ref: Cyclosporine Test to detect cyclosporin levels. This test can be performed on Bovine, Canine, Equine, Feline, Ferrets, Fruit Bats, Ovine, Porcine, Primate samples only. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	66.00 (+ referral fee; see comment)	Referral	W, F	3-5 business	1 mL EDTA whole blood	Lavender top blood collection tube	Refrigerate; DO NOT FREEZE. Ship for next day delivery in insulated container with freezer pack(s)
Ref: Digoxin Checking Digoxin levels in the blood Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Do not store in rubber topped tubes as digoxin will bind to the rubber. Fasting is recommended, but not required.	60.00 (+ referral fee; see comment)	Referral	Run when received	2-4 business	1 mL serum	red top blood collection tube; GLASS vial	Refrigerate. Ship in insulated container with ice pack.
Ref: FISH Fluorescent In-Situ Hybridization Fluorescent probe binds to bacterial ribosomes in tissue. Detected with fluorescent microscope. This test can be performed on all species. A \$15 handling fee will be added to each sample, so the total price for a single tissue is \$190. Additional tissues will be billed at \$50 for each additional tissue. Additional probes will be billed at \$50.00 for each additional probe. For specific information regarding samples, instructions and forms, please see FISH Fluorescent In-Situ Hybridization. Research questions should be directed directly to Simpson Laboratory (607-253-3567).	175.00 + in-house referral fee. See comment	Referral	Tues or Wed	5-10 Business	5 unstained fixed tissue slides.	Slide Mailer;	Room Temperature
Ref: Insulin Growth Factor-1 Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Most species can be tested.	75.60 (+ referral fee; see comment)	Referral	Wednesday	4 - 9 business	0.5 mL Serum or Plasma	Red Top or Purple Top, SST	Refrigerate. Ship in insulated container with cold pack
Ref: Keppra/Levetiracetam Checking Keppra levels in the blood Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Fasting is recommended, but not required.	84.00 (+ referral fee; see comment)	Referral	Tues, Thurs	3-4 business	1 mL serum	Red top blood collection tube	Refrigerate. Ship in insulated container with ice pack
Ref: Lawsonia IPMA Referral test prices are set by the referral laboratory and may change without notice. a \$30.00 handling fee is added for each accession sent to each referring lab	24.00 (+ referral fee; see comment)	Referral	M-F	7-10 days	1 mL serum		Refrigerate

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Ref: Leishmania PCR Sample Required is EDTA whole blood. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	60.00 (+ referral fee; see comment)	Referral	M-F	5-7 days	2 mL EDTA whole blood	Insulated	Refrigerate or ship on cold pack
Ref: Malignant Catarrhal Fever IPT This is an immunoperoxidase test designed to detect antibodies to the MCF virus. This serology test is indicated for clinical suspects. This test cross reacts with both Alcelaphine-associated and sheep-associated virus antibodies. All antibody positive IPT samples will be tested by the VN, resulting in two charges (IPT and VN). A clinical suspect is expected to be positive on the IPT test and negative on the VN test. Please call the laboratory before sending sample. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	20.40 (+ referral fee; see comment)	Referral	Wednesday	5-7 days	2 mL serum	red top tube	Refrigerate. Ship in insulated container with ice packs
Ref: MCF PCR PCR test to detect the Herpes viruses recognized to cause Malignant Catarrhal Fever. This PCR assay differentiates between alcelaphine herpesvirus 1 and ovine herpesvirus 2 The sample required from a live animal is EDTA whole blood. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	67.20 (+ referral fee; see comment)	Referral	M-F	7-14 days	EDTA Whole Blood or Tissue	Insulated	Refrigerate (not frozen)
Ref: Mycobacterial Culture (Acid Fast TB) Test for presence of tuberculosis. This test does not speciate non-TB Mycobacterium species Send via USPO in NVSL mailer. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. This test is intended to detect Mycobacterium tuberculosis as a reportable disease.	58.00 (+ referral fee; see comment)	Referral	M-F	30-120 days	Tissue	Mycobacterium box	Ambient
Ref: Mycobacterial PCR PCR for Mycobacterium species. Not for M. bovis or M. tuberculosis suspects. Formalin fixed tissue or paraffin blocks should not be decalcified. Tissue should be in wet formalin less than 10 days. Acid fast bacteria must be present on histopathology to perform PCR testing. Fresh tissue should be tested by requesting Mycobacterium Culture rather than PCR to increase testing sensitivity.	246.00 (+ referral fee; see comment)	Referral	M-F	12-14 Busine:	Fixed tissue containing lesion	Sterile, leakproof, unbreakable container or paraffin block	Room temperature for fixed tissue and paraffin block
Ref: Mycobacterium Culture ID All culture isolates (positive cultures for mycobacterium) will require additional speciation at a fee of \$70.00 to identify the Mycobacteria further. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	48.00 (+ referral fee; see comment)	Referral	Mon - Fri	60 - 120 days	Fresh tissue	Sterile, leakproof, unbreakable container	Ship for next day delivery in an insulated container with freezer packs.
Ref: Mycopl hyopneumoniae PCR This test detects nucleic acid of the Mycoplasma hyopneumoniae bacteria Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	48.36 (+ referral fee; see comment)	Referral	M - F	1 - 3 days	lung, bronchial swab, BAL	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs
Ref: Mycopl hyorhinis PCR This test detects the nucleic acid of the Mycoplasma hyorhinis bacteria Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	30.00 (+ referral fee; see comment)	Referral	M, Th	3 - 5 days	joint swab or joint tissue	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs.
Ref: Mycopl hyosynoviae PCR This test detects the nucleic acid of the Mycoplasma hyosynoviae bacteria Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	30.00 (+ referral fee; see comment)	Referral	M, Th	3-5 days	joint swab or joint tissue	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Ref: Parathyroid Hormone Used in diagnosis of: Calcium metabolic disorders, Hypoparathyroidism, Hyperparathyroidism – Primary or Secondary, Malignancy, Vitamin D disorders, Renal disease, Hypercalcemia, Hypocalcemia. Also used to monitor calcitriol therapy. This test can be performed on samples from all species. Send via overnight courier. Avoid Hemolysis and lipemia. Fast animal overnight before taking sample. Ionized Calcium is automatically added to all Parathyroid Hormone requests at no additional charge. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.	57.60 (+ referral fee; see comment)	Referral	M-F	2-6 Business	1 mL serum only	Red top blood collection tubetube	Refrigerate. Ship for next day delivery in insulated container with freezer pack(s)
Ref: Por E coli Genotyping PCR Detects the genes responsible for producing the following pathogenic markers: STb (toxin), Sta (toxin), K99 (pilus), LT (toxin), F18 (pilus), K88 (pilus), F41 (pilus), Stx2e (toxin), EAST1 (toxin), Stx 1 (toxin), Stx 2 (toxin), AIDA (adhesin), PAA (adhesin) Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	36.00 (+ referral fee; see comment)	Referral	M, Th	5 - 7 days	pure culture of isolate	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs
Ref: Por Epidemic Diarrhea PCR This test detects the nucleic acid of the porcine epidemic diarrhea (PED) virus. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	36.00 (+ referral fee; see comment)	Referral	M - F	1 - 3 days	feces, fecal swab, intestine, oral fluids, environmental samples	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs.
Ref: Porc cytomegalovirus PCR Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	42.00 (+ referral fee; see comment)	Referral	M, Th	3 - 5 days	blood, nasal swab, nasal turbinates, brain	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs.
Ref: Porc deltacoronavirus PCR This test detects the nucleic acid of the porcine delta coronavirus. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	36.00 (+ referral fee; see comment)	Referral	M - F	1 - 3 days	feces, fecal swab, intestine, oral fluids, environmental samples	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs
Ref: Porcine Circovirus IFA Iowa State charges a \$12.00 submission fee. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring labSerum samples are stable at room temperature for several days. If longer, refrigerate or freeze.	21.00 (+ referral fee; see comment)	Referral	M, F	3-5 days	2 mL serum	Red top blood collection tube	It is recommended to ship samples with an ice pack.
Ref: Porcine Parvovirus HI This test detects antibodies to the porcine parvovirus Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	6.00 (+ referral fee; see comment)	Referral	Mondays	1 - 3 days	serum	red top tube	Ship in insulated container with ice packs
Ref: Porcine PED/TGE PCR This test detects the nucleic acid of the porcine epidemic diarrhea (PED) virus and the transmissible gastroenteritis (TGE) virus. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	42.00 (+ referral fee; see comment)	Referral	M - F	1- 3 days	feces, fecal swab, intestine, oral fluids, environmental samples	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs.
Ref: Pseudorabies gB ELISA Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	16.00 (+ referral fee; see comment)	Referral	3 X per week	2-4 days	1 mL Serum	Red top tube	Refrigerate
Ref: Pseudorabies PCR This test detects the nucleic acid of Pseudorabies virus Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	36.00 (+ referral fee; see comment)	Referral	T, F	5 - 8 days	brain, tonsil, lung, fetal liver	unbreakable, leak-proof container	Refrigerate
Ref: Quantitative Calculi/Urinary This test provides an analysis of the crystalline composition of the urolith(urinary stone) Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab	38.00 (+ referral fee; see comment)	Referral	M-F	4-6 business	Stone	Baylor Stone Collection Cup	None

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Ref: Rabies Titer	83.00 (+ referral fee; see comment)	Referral	M, Th	23-30 Busine:	2 mL serum	Red top blood collection tube	Refrigerate. Ship for next day delivery in insulated container with freezer pack(s).
<p>This test is not used for export testing. This test provides an endpoint titer. Please see comment above regarding export testing. This test can be performed on all species. Send via overnight courier. Can use heat inactivated serum, must be noted on the form. Cannot use plasma. Sample needs at least 500ul for one run. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. The AHDC does not refer Rabies titers for export testing due to requirements of the testing lab. Please contact Kansas State Veterinary Diagnostic Lab directly at 785-532-4483.</p>							
Ref: Stone Analysis - Minnesota	0.00 (+ referral fee; see comment)	Referral	M-F	10-14 days	Stone	Unbreakable container	
<p>Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab</p>							
Ref: TGE/Por Resp Corona ELISA	11.00 (+ referral fee; see comment)	Referral	T, Th	1 - 3 days	serum	unbreakable, leak-proof container	Refrigerate. Ship in insulated container with ice packs
<p>This test detects antibodies to the transmissible gastroenteritis (TGE) virus/ porcine respiratory coronavirus. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab</p>							
Ref: Theophylline	60.00 (+ referral fee; see comment)	Referral	Run as received	2-4 business	1 mL serum	red top blood collection tube; glass or plastic vial	Refrigerate. Ship in insulated container with ice pack
<p>Checking Theophylline levels in the blood Send via overnight courier Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab. Please do not use Serum Separator Tubes. Fasting is recommended, but not required</p>							
Ref: Tick Panel - PCR	150.00 (+ referral fee; see comment)	Referral	Batched weekly	9-12 business	2 mL EDTA whole blood	EDTA blood Tube	Refrigerate, ship for next day delivery in insulated container with freezer pack(s)
<p>PCR Panel includes: PCR testing for known pathogens in six genera: Babesia, Bartonella, Anaplasma, Ehrlichia, Rickettsia, and hemotropic Mycoplasma. All positive PCR results are verified by performing sequencing analysis which also takes identification to the species and sometimes strain level. This test can be performed on most species. Call lab with exotic animals. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p>							
Ref: Troponin	88.00 (+ referral fee; see comment)	Referral	M, F	1-3 days	1 mL serum or hep plasma	Insulated	Dry ice
<p>A complex of muscle proteins which, when combined with Ca²⁺, influence tropomyosin to initiate contraction. Send pre and post exercise samples on dry ice by overnight courier. Remove serum or plasma from cells within 2 hours! Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab</p>							
Ref: Vitamin D Profile	76.80 (+ referral fee; see comment)	Referral	Monday and Friday	2-8 business	1.5 mL Serum or Heparinized plasma(not EDTA plasma)	Red Top, Purple Top, Green Top or SST blood tubes	Refrigerate, ship for next day delivery in insulated container with freezer pack(s)
<p>Assay for 25- Hydroxyvitamin D This test can be performed on most species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p>							
Ref: Vitamin D, 25-hyd	56.40 (+ referral fee; see comment)	Referral	Monday and Friday	3-9 Business	0.5 mL Serum or Plasma (Heparinized and EDTA is accepted)	Red Top, Purple Top, Green Top or SST blood tubes	Red Top, Purple Top, Green Top or SST blood collection tubes
<p>1. To identify Vitamin D toxicity or deficiency. A. Will pick up metabolites of cholecalciferol (rat poisons, vitamin D supplements). B. Will not pick up Vitamin D toxicity due to calcitriol (Rocaltrol) or calcipotriene (Dovonex - anti-psoriasis cream). 2. To identify Vitamin D deficiency in animals eating homemade diets, or for certain species, especially Llamas/Alpacas, lizards, primates, and other zoo animals ?This test can be performed on appropriate samples from all species. Send via overnight courier. Referral test fee is set by the referring lab and may change without notice; also, a \$30.00 handling fee is added for each accession sent to each referring lab.</p>							
Reserpine (LC/MS)	165.50 /sample	Toxicology	M-F	1-3 days	1) 10 mL heparin whole blood; or 2) transfer 5 mL heparin plasma to a red top tube	1) Heparinized tube, no gel clot; 2) red top blood collection tube	1-2) Refrigerate w/ cool pack
<p>Liquid Chromatography/Mass Spectrometry (LC/MS) (increased sensitivity)</p>							
Reticulocyte Count	10.50	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate
<p>Do not allow coolant to come in contact with specimen; ship Overnight</p>							

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Rivaroxaban Anti-Xa activity measured in a chromogenic assay configured with a rivaroxaban calibration standard. Samples for determination of peak rivaroxaban anti-Xa activity should be collected 3 hours after oral drug administration. For more information, see Rivaroxaban assay FAQs	45.00	Comparative Coagulation	As required	Same day	0.5 mL Citrated plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate sample. Ship overnight with ice packs in insulated container.
Rotavirus FA-Any Species Fluorescent antibody	25.00	Virology	M - F	2 - 4 days		Sealable plastic bag or container	Ship in insulated container with ice packs.
Salmonella Culture For the isolation of Salmonella species. *May qualify for New York State contract pricing. Test is also part of the Enteric Panel 1, Enteric Panel 2, and Enteric Panel 3. Antimicrobial susceptibilities are not included in the testing fee. See the Adult Equine Diarrhea Kit for equine submissions.	22.00 /sample	Bacteriology	M-Sa	3-5 days	1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) other tissues; 5) Water; 6) Swab of 1-4	1) Feces in sealed sterile container or ParaPak; 2-5) Tissues or fluids in separate, sealed, sterile containers 6) Swab in Amies transport media or other suitable bacterial transport medium	1-6) Refrigerate EXCEPT ParaPak (Parapak at room temperature)
Selenium, tissue/feed (Selenium Hydride Atomic Absorption)	55.00	Toxicology	M-F	6 days	1) 2g tissue; 2) 100g feed	1) sterile container; 2) any container	1) Refrigerate; 2) none
Selenium, whole blood Atomic Absorption analysis of whole blood Cannot analyze in plasma or serum - Need whole blood. Contact Lab for specifics for avian and very small animals. May also want to consider "Vitamin E Alcohol in Serum" test as well.	28.00 /sample	Toxicology	M-F	3 days	1) 5 mL heparin whole blood; 2) 5 mL EDTA whole blood	1) green top blood collection tube; 2) purple top blood collection tube	1-2) Refrigerate w/ cool pack
Skin Scraping For ectoparasites Visual examination, KOH digest and Sugar Centrifugation Flotation; send an adequate sample size in an escape-proof container; remember to scrape until skin begins to bleed	28.00	Parasitology	M-F	1-3 days	hair/fur and epidermal tissue	escape-proof, sealed container	Not needed
Sodium Sodium, urine	9.00 /sample (blood, fluid) 17.00	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid 1 mL urine	plain glass or plastic tube plain glass or plastic tube	Refrigerate Refrigerate
Sorbitol Dehydrogenase (SDH), blood	11.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Special Stain #1 THIS TEST CODE FOR INTERNAL USE ONLY.	50.00	Clinical Pathology	M - F	5 days	Slides		
Special Stain #2	80.00	Clinical Pathology	M-F	5 days	Slides		
Special Stain #3	110.00	Clinical Pathology	M - F	5 days	SLIDES		
Special Stain #4	140.00	Clinical Pathology	M - F	5 days	Slides		
Special/Augmented Biopsy Surcharge Intended for complex submissions of entire organs, lobes, systems, tumor beds, limbs, jaws, etc. This includes research/insurance/legal cases for which every single tissue submitted must be described histologically and the selection of the tissues is not at the discretion of the pathologist. It may also include complicated submissions of entire organs or tumor beds or lymphatic samples that need exact tracking of site of collection for multiple samples of the same tissue type. Please call 607-253-3319 in advance to discuss research histology needs or for price quotes related to these types of submissions.	56.00	Anatomic Pathology	Mon - Fri	4-10 days	0.5 to 1.0 cm thick section of tissue fixed in 10% neutralized formalin, 10:1 formalin to tissue ratio.	Wide-mouth, leakproof, unbreakable jar; NOT GLASS	Store at room temperature. Do not freeze

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Stained Histology Slides Slides prepared from (FFPE) blocks H&E or Special Stained slides prepared from Anatomic Pathology cases in addition to the Histopathology. Stained slides are requested to either have Anatomic Pathology review or to have them sent to another facility for review/referral. Additional charges will be applied for shipping and handling.	10.25	Anatomic Pathology	M-F	5 business days	Paraffin embedded tissue block		Room temperature
Strychnine, liquids/fluids	55.25 /sample	Toxicology	M-F	3 days	1) 10 - 20 mL heparin whole blood; 2) 10 - 20 mL urine	1) Heparinized tube, no gel clot; 2) sterile leak-proof container	1-2) Refrigerate w/ cool pack
Strychnine, solids	90.00 /sample	Toxicology	M-F	3 days	1) 50g liver; 2) 50g stomach contents or vomitus	1-2) sterile container	1-2) Refrigerate w/ cool pack
Sulfur	40.00	Toxicology	M-F	3 days	1) 2g tissue; 2) 100g feed	1) sterile container; 2) any container	1) Refrigerate; 2) none
Swine Coronavirus (TGE) PCR The alpha coronavirus PCR employed by the AHDC will detect feline enteric coronavirus, canine enteric coronavirus, ferret or mink coronavirus, and swine coronavirus - AKA transmissible gastroenteritis virus. For species specific testing information please see Coronavirus PCR Guidelines.	36.75	Molecular Diagnostics	M-F as needed	3-5 days	1) tissue (min. 5 g); 2) ascites or body fluid (min. 2 mL); 3) feces (min 5 g)	unbreakable, leak-proof container	Refrigerate
Synovial Fluid Cytology Report includes - Specimen type and site (provided by submitter), Volume, Color, Turbidity, Total Protein, Nucleated cell count, RBC count, Viscosity, Smear type, Smear description, Smear Interpretation If only slides (and no fluid) are submitted, request a cytology smear exam on the cytology submission form and specify the joint sampled as the source. For joint fluid in EDTA, please also submit 2-3 freshly prepared smears from the fluid. Also specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. It is important to rapidly air-dry any smears. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing	55.00	Clinical Pathology	M - F	1 business day	1) Joint fluid in EDTA AND 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only; ship in insulated container with freezer pack for next day delivery. Hold smears at room temperature; protect from chilling/coolant by insulating separately during shipment.
Testosterone baseline For more information, see Appendix C	19.00	Endocrinology	Monday	1-8 days	1 mL serum	plastic or glass tube	Refrigerate
Testosterone Post HCG/GnRH	13.00	Endocrinology	Monday	1-7 days	250 uL serum or plasma	plastic or glass	
Testosterone: HCG/GnRH Response	19.00	Endocrinology	Monday	1-7 days	250 uL serum or plasma	plastic or glass	
Thrombin-antithrombin Complex Research and batch testing only. Testing not done on rabbits. Per test fee requires submission of 15 or more samples for batch analysis. Ship Overnight. For additional info on Submission Guidelines, or see Comp Coag web page;	52.50	Comparative Coagulation	1x/month	up to 1 month	citrate plasma	plastic tube in styrofoam box	dry ice
Total Protein Panel Total Protein (TP), Albumin, Globulin for additional info on Clin Path panels	17.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Total Protein Plasma (by refractometer) Do not allow coolant to come in contact with specimen; ship Overnight	5.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (>1/2 full)	lavender top tube	Refrigerate
Total Protein, blood Total protein in blood	13.00	Clinical Pathology	M-Sa	1-2 days	1 mL separated serum or heparinized plasma	plain glass or plastic tube	Refrigerate
Total Protein-Creatinine Panel Total Protein, Creatinine, P:C Ratio (Calc) for additional info on Clin Path panels	38.00	Clinical Pathology	M-Sa	1-2 days	2 mL urine	leak-proof container	Refrigerate

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 27 of 30

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Toxic Heavy Metal Screen (Arsenic, Lead, Cadmium, Mercury, Thallium) X-ray Fluorescence Spectrophotometry	77.25 /sample	Toxicology	M-F	7 days	1) liver OR 2) kidney OR 3) powders, solids OR 4) paint chips	1-5) sterile, leak-proof container	1-2) Freeze; 3-5) Refrigerate w/ cool pack
Toxoplasma gondii IgG ELISA Detects anti-Toxoplasma IgG antibody The Toxoplasma gondii IgG ELISA tests serum for the presence of anti-Toxoplasma IgG antibody, indicating a history of exposure to the parasite. This test is valid to test serum from a broad range of animal species, except for birds, marsupials and cold-blooded animals. Results are reported as "positive", "negative" or "equivocal". Testing is repeated 2-3 weeks after the initial sample was tested to establish if an infection is acute or chronic, or to delineate the true status of an animal with a previously "equivocal" result. When a second sample is submitted, please provide the accession number from the first sample submitted, as both samples are needed for accurate paired testing. Fluids other than serum will not be evaluated. *May qualify for New York State contract pricing.	20.00 /sample price.	Parasitology	Wed	1-7 days	100 uL serum	red-top blood collection tube	Refrigerate
Toxoplasmosis FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fetal tissue: placenta and brain preferred; also lung, liver, kidney, skeletal muscle, spleen, lymph node; Non-fetal tissue including suspect affected tissue such as brain, liver, kidney, lung, skeletal muscle.	Leakproof, unbreakable plastic container or bag	Ship for next day delivery in insulated container with freezer pack(s)
Tracheal Wash Cytology Report includes - Specimen type and site (provided by submitter), Smear type, Smear description, Smear Interpretation With fluid samples of tracheal wash, please also submit 2-3 freshly prepared smears of the fluid and specify smear type, i.e. direct or sediment, on the smears and on the cytology submission form. It is important to rapidly air-dry any smears. Do not allow coolant to come in contact with specimen; ship overnight. STAT service available for same day results (extra charges apply). For additional information see Cytology Testing	46.00	Clinical Pathology	M - F	1 business day	1) Tracheal wash fluid in EDTA; 2) Smears (at least 2-3 unstained smears preferred); 3) Also submit a red-top tube if culture is anticipated.	1) Lavender-top tube; (+/- red-top tube); 2) Frosted-end slides	Refrigerate fluids only, DO NOT refrigerate smears
Transition Cow Energy Panel Includes BHB and NEFA	22.00	Clinical Pathology	M-Sa	4 hrs	1 mL separated serum	plain glass or plastic tube	Refrigerate
Transmissible Gastroenteritis (TGE) Virus FA Fluorescent antibody	25.00 /animal	Virology	M-F	2-4 days	Fresh Ileum, jejunum, cecum, lung	Sealable plastic bag or container	Ship in insulated container with ice pack
Trichinella Tissue Examination Acid Pepsin digest	27.50	Parasitology	M-F	7 days	muscle tissue	wide mouth leak-proof	Refrigerate
Trichomonas Fecal Examination Trichomonas pouches are available from AHDC Shipping	25.00	Parasitology	M-F	12 days	fecal swab in trichomonas In-Pouch (see instructions with In-Pouch)	trichomonas In-Pouch	Do NOT Refrigerate
Triglycerides	11.00 /sample (blood, fluid)	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate
Unstained Histology Slides Prepared Slides or PCR tubes prepared from (FFPE) blocks. Unstained slides or PCR tubes containing 2-3 (10uM) unstained tissue sections prepared from anatomic pathology cases that have been or are in the process of being reported on. These unstained slides or PCR tubes are prepared for the purpose of sending to another testing facility for slide review or referral analysis. Additional charges may be applied for shipping and handling. Price is per slide or PCR tube prepared.	5.25	Anatomic Pathology	M-F	5 business days	Paraffin embedded tissue block		Room temperature
Urea Nitrogen Test for blood urea nitrogen in serum, plasma or other fluid.	11.00	Clinical Pathology	M-Sa	1-2 days	1) 1 mL separated serum or heparinized plasma; 2) fluid	plain glass or plastic tube	Refrigerate. Ship in insulated container with ice pack.
Urea Nitrogen, urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate. Ship in insulated container with ice pack.
Ureaplasma Culture For the culture of Ureaplasma species. *May qualify for New York State contract pricing. Antimicrobial susceptibilities are not applicable. Test is also part of the Reproductive Panel 4 and Respiratory Panel 6.	27.00 /sample	Bacteriology	M-F	5-10 days	1) Uterine, cervical or vaginal mucus; 2) Semen; 3) Tracheal washing or lung tissue; 4) Urine; 5) Aseptically collected swab of 1-4	1-4) Fluid or tissue in sealed sterile container; 5) Swab in Amies transport media WITHOUT charcoal	1-4) Refrigerate

Last Updated: 10/24/2017 * Each accession received will be charged a \$5.00 accessioning fee, in addition to the requested testing. Fees are subject to change without notice, please call the lab if you have any questions. Contract pricing available. For information please refer to <https://ahdc.vet.cornell.edu/billing/contract.cfm>
VSS-WEB-084-V01 ** "Lag Time" is counted in business days, unless otherwise specified, from the day the specimen arrives until the day the test result is reported, assuming there are no complications with testing.

Page 28 of 30

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
Uric Acid, urine	16.50	Clinical Pathology	M-Sa	1-2 days	1 mL urine	plain glass or plastic tube	Refrigerate
Urinalysis, Routine Color, turbidity, dipstick (pH, TP, glucose, ketones, bilirubin, blood), specific gravity, sediment exam	25.00	Clinical Pathology	M-Sa	1-2 days	10 mL fresh urine	leakproof container	Refrigerate
Urine Culture For the isolation of aerobic bacteria from the urinary tract. *May qualify for New York State contract pricing. It is recommended to collect and submit urine collected by Cystocentesis. Please indicate method of collection. Antimicrobial susceptibilities are not included in the testing fee. Refer to Urine Antimicrobial susceptibilities. Urine Transport Tubes are available for purchase from our Supply Center. Performing cytology in conjunction with culture is also recommended to aid in interpretation of results.	20.00 /sample	Bacteriology	M-F	2-7 days	1) Urine; 2) Bladder tissue/stones; 3) Aseptically collected Swab of urine or bladder	1) Urine Transport Tube (preferred) or other sterile container; 2) Tissue or stones in separate sterile sealed containers; 3) Swab in Amies transport media or other suitable transport medium for aerobic culture	(1-3) Ship overnight in insulated container with ice packs to arrive no later than 48 hours post collection.
Vesicular Stomatitis Virus (VSV) IND SN IND strain Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation.	10.00	Virology	M, W, F	2-4 days	1 ml serum	red top blood collection tube	It is recommended to ship samples with an ice pack.
Vesicular Stomatitis Virus (VSV) NJ SN NJ strain Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Paired serum samples are recommended for optimal diagnostic test interpretation.	10.00	Virology	2-4 days		1 ml serum	red top blood collection tube	It is recommended to ship samples with an ice pack.
Virus FA See also specific Viral FA tests Refer to individual FA tests	25.00 /animal	Virology	M-F	2-4 days	Tissues or fluids. Contact lab if there are questions about sample suitability.	Sealable plastic bag or container for tissues; Leakproof container for fluids and swabs in Viral Transport Medium	Ship in insulated container with ice pack
Virus Isolation See also specific Virus Isolation tests or clinical presentation Best sample may depend upon suspected viral infection and pathophysiology related to viremia, infected tissue and viral shedding. Recent investigations indicate that urine is a good antemortem sample for detection of some viruses. There is generally a lack of information regarding viral detection in urine samples for many viral diseases. Please include urine if it is available when you are submitting other samples for virus isolation for our investigational use. No extra charges will be incurred. Refer also to individual Virus Isolation tests. Call lab ahead before sending reptile samples. Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. If viral transport media is unavailable, put swabs in a red top tube with a few drops of saline.	70.00 /animal	Virology	M-F	21-30 days, s	1) Tissue [target organs and blood-rich organs such as lymph nodes, spleen, lungs, kidney, heart, liver, brain] OR 2) Cerebrospinal fluid Or 3) EDTA whole blood OR 4) antemortem nasal swab AND 5) urine (see comment)	1) Tissue in leakproof unbreakable containers 2) sterile, leakproof container; 3) EDTA blood tube; 4) in sterile leakproof container with 2-3 drops sterile saline or in Virus isolation media (not bacterial transport media)	Ship in insulated container with freezer packs for next day delivery pack
Vitamin A in serum (High Performance Liquid Chromatography) Analysis for retinol (vitamin A) using HPLC. Separate serum and remove from clot immediately. Protect from light and keep cool.	30.00 each	Toxicology	Tu, Th	5 days	Serum, separated from clot.	Avoid preservatives such as Citrate or EDTA. No Gel Clot Vacutainer Tubes.	Refrigerate
Vitamin E in serum (High Performance Liquid Chromatography) "Selenium" test recommended also Analysis for alpha-tocopherol (vitamin E) using HPLC. Separate serum and remove from clot immediately. Failure to remove serum from clot may lead to inaccurately low vitamin E levels. Protect from light and keep cool. We strongly recommend that Vit E (serum) and Selenium (Heparin or EDTA whole blood) be tested together.	30.00 each	Toxicology	Tu, Th	5 days	Serum, separated from clot	Avoid preservatives such as Citrate or EDTA. No Gel Clot Vacutainer Tubes (SST tubes).	Refrigerate

Animal Health Diagnostic Center Test and Fee Schedule

Test Name	Test Fee	Discipline	Test Days	Lag**	Samples	Container	Coolant
von Willebrand factor - large animal/exotic von Willebrand factor concentration, ELISA method Test offered for species other than dog and cat. For additional info go to Submission Guidelines, or see Comp Coag web page	48.50	Comparative Coagulation	1x/wk	14 days	1 mL citrate plasma (separated) EDTA plasma is acceptable	plastic or siliconized glass tube in styrofoam box	Cold pack or dry ice
von Willebrand factor Collagen Binding Assay Functional ELISA Ship Overnight; requires vWF:Ag on same sample; for additional info on Submission Guidelines, or see Comp Coag web page	43.00	Comparative Coagulation	bi-weekly	1-2 wks	citrate plasma	plastic or siliconized glass tube in styrofoam box	Refrigerate/ship on cold pack
West Nile Virus (WNV) PCR PCR Swabs in bacterial transport medium are not appropriate for virus isolation or PCR testing. Whole blood is an acceptable sample only for Avian testing, not for Mammals This service is performed pursuant to an agreement with Roche Molecular Systems.	36.75	Virology	M-F	3-5 days	AVIAN: 1) Brain, heart, kidney, bone marrow; 2) EDTA Whole Blood; 3) Oropharyngeal or Cloacal swab; MAMMAL: 4) CSF; 5) Brain or spinal cord. WHOLE BLOOD IS NOT AN ACCEPTABLE SPECIMEN FOR MAMMALS	1,5) Sealable plastic bags or containers; 3) Swabs in Virus transport medium; 2,4) Leakproof container	Freezer Packs
West Nile Virus (WNV) SN serum neutralization Serum samples are stable at room temperature for several days. If longer, refrigerate or freeze. Acute and convalescent serum samples are recommended for optimal diagnostic test interpretation. Currently at the AHDC, this test is used primarily for mammalian species.	14.50	Virology	Thursday	7-14 days	1 ml of serum (preferred) or plasma	red top blood collection tube	It is recommended to ship samples with an ice pack.
White Blood Cell (WBC) Count, Total Do not allow coolant to come in contact with specimen; ship Overnight	7.00	Clinical Pathology	M-Sa	1-2 days	EDTA blood (> 1/2 full)	lavender top tube	Refrigerate
White Blood Cell (WBC) Panel WBC, Manual Differential Cell Count, Platelet Smear estimate, RBC, WBC Do not allow coolant to come in contact with specimen or slides; ship overnight. For additional info on Clin Path panels	16.50	Clinical Pathology	M-Sa	1-2 days	1) EDTA blood > 1/2 full; 2) 2 dry (unfixed, unstained) blood smears	1) lavender top tube; 2) frosted-end slides	Refrigerate. Ship in insulated container with ice packs.
Yersinia Culture For the isolation of Yersinia enterocolitica. Antimicrobial susceptibilities are not included in the testing fee.	16.00 /sample	Bacteriology	M-Sa	2-7 days	1) Fresh feces or intestinal contents; 2) Ligated ileum or jejunum; 3) Mesenteric lymph node; 4) swab of 1-3	1) Feces in sealed sterile container or ParaPak; 2-3) Tissue in Separate sealed sterile containers; 4) Swab in Amies transport media or other suitable aerobic transport medium	1-3) Refrigerate EXCEPT ParaPak (keep at room temperature)
Zinc, serum/plasma For avian and exotics, contact Lab	30.00 /sample	Toxicology	M-F	2 days	1) 2 mL serum; or 2) 2 mL heparinized plasma in a green top tube	1) 2 mL serum; or 2) 2 mL heparinized plasma; Royal blue-topped tube (trace mineral tube) preferred. No gel clot.	1-2) Refrigerate. Ship in insulated container with ice pack.